Quick Start Guide

GenElute™ Gel Extraction Kit

NA1111

Reagents to Prepare

 Dilute the Wash Solution Concentrate G with 48 mL of 95-100% ethanol. After each use, tightly cap the diluted Wash Solution to prevent ethanol evaporation.

Protocol

All spins at $12,000-16,000 \times g$.

Sample Preparation

- 1. Excise the DNA fragment of interest and weigh the gel slice in a colorless tube.
- 2. Add 3 gel volumes of Gel Solubilization Solution and incubate the mixture at 50-60 °C for 10 min or until the gel slice is completely dissolved. Vortex every 2-3 min during incubation to dissolve the gel.
- 3. Once gel slice is dissolved ensure the color of the mixture is yellow. If the mixture color is red, add 10 μ L of 3M Sodium Acetate, pH 5.2 and mix. Continue adding 3M Sodium Acetate, pH 5.2 in increments of 10 μ L until the mixture turns yellow.
- 4. Add 1 gel volume of 100% isopropanol and mix until homogenous. If the agarose concentration is > 2%, use 2 gel volumes of 100% isopropanol.

Bind DNA

- 5. Place Column into a 2 mL collection tube.
- 6. Add 0.5 mL Column Preparation Solution to each column and spin for 1 min. Discard the flowthrough.
- 7. Load 700 μ L solubilized gel mixture into column and spin for 1 min. Discard flowthrough. Repeat if the volume is > 700 μ L.

Wash to Remove Contaminants

- 8. Add 700 μL of Wash Solution to the column and spin for 1 min. Discard the eluate.
- Replace column with its collection tube and spin for an additional 1 min to remove excess ethanol. Discard the eluate and collection tube.

DNA Elution

- 10. Transfer column to fresh collection tube.
- 11. Add 50 μ L Elution Solution (pre-heated to 65 °C) to the center of column. Incubate at room temperature for 1 min.
- 12. Spin column for 1 min.

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/Offices.

Technical Assistance

Visit the tech service page on our web site at SigmaAldrich.com/TechService.

Standard Warranty

The applicable warranty for the products listed in this publication may be found at SigmaAldrich.com/Terms.

Merck, Millipore, GenElute and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

 $\ \, \odot$ 2022 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.



