## **Quick Start Guide**

# GenElute™ HP Endotoxin-Free Plasmid Maxiprep Kit

## NA0300S, NA0300, NA0310

## Reagents to Prepare

- Spin RNase A Solution tube briefly and add 750 μL (4 and 10 preps) or 1.9 mL (25 preps) of RNase A Solution to the Resuspension Solution. Store at 4 °C.
- Add 120 mL (4 and 10 preps) or 300 mL (25 preps) of 95-100% ethanol to Wash Solution 2. After each use, tightly cap the diluted Wash Solution to prevent ethanol evaporation.

## **Protocol**

Vacuum format.

#### **Harvest Bacteria**

 Pellet cells from 150 mL overnight culture by spinning for 10 min at 5,000 x q. Discard supernatant.

## **Resuspend and Lyse Bacteria**

- Resuspend cells in 12 mL Resuspension Solution. Pipet or vortex to mix.
- 3. Add 12 mL Lysis Solution. Gently invert 6-8 times to mix. Do not vortex. Allow to clear for 3-5 min.

## **Prepare Cleared Lysate**

- 4. Remove plunger from Filter Syringe and place the barrel in an upright position.
- 5. Add 12 mL Neutralization Solution (chilled prior to use) to lysed cells and gently invert 6-8 times.
- 6. Add 9 mL Binding Solution and gently invert 1-2 times. Immediately add the mixture to the barrel of the filter syringe and let sit for 5 min.

## **Prepare Column**

- 7. Place Column onto vacuum manifold and apply vacuum.
- 8. Add 4 mL Column Preparation Solution to the column and allow it to pass through.

## **Bind DNA to Column**

9. Hold Filter Syringe over the column and gently insert the plunger to expel the cleared lysate. Allow the lysate to pass through the column.

### **Wash to Remove Contaminants**

- 10. Add 12 mL Wash Solution 1 to column and allow it to pass through.
- 11. Add 12 mL Wash Solution 2 to column and allow it to pass through.
- 12. Maintain vacuum until the column is dry (10 min). If more than 6 columns are on the manifold, dry for at least 20 min.

## **Elute Purified DNA**

- 13. Transfer column to a collection tube.
- 14. Add 3 mL Elution Solution to the column.
- 15. Spin in a swinging bucket rotor at 3,000  $\times$  g for 5 min for maximum plasmid DNA yield or 1,000  $\times$  g for 5 min for maximum plasmid DNA concentration.

## **Notice**

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

### **Contact Information**

For the location of the office nearest you, go to <u>SigmaAldrich.com/Offices</u>.

### Technical Assistance

Visit the tech service page on our web site at SigmaAldrich.com/TechService.

## **Standard Warranty**

The applicable warranty for the products listed in this publication may be found at <u>SigmaAldrich.com/Terms</u>.

MilliporeSigma, Millipore, GenElute and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2022 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the U.S. and Canada.

