

Product Information

Anti-Fibronectin antibody, Mouse Monoclonal

Clone IST-4, Purified from Hybridoma Cell Culture

F0916

Product Description

Anti-Fibronectin antibody, Mouse Monoclonal is derived from the IST-4 hybridoma produced by the fusion of mouse myeloma cells and splenocytes from immunized BALB/c mice. Purified human plasma fibronectin was used as the immunogen.^{1,2}.

Anti-Fibronectin antibody (clone IST-4) recognizes an epitope located within the 5th type III repeat of human plasma fibronectin. 1,3-5 This epitope is common to all fibronectin forms. 4,6 The antibody localizes plasma and cellular fibronectin both in its native and denatured-reduced forms (240 kDa) applying immunoblotting, RIA, ELISA, immunocytochemistry and immunohistochemistry. It may be used for the specific labeling of fibronectin in cultured cells and in frozen tissue sections. Staining of formalin-fixed, paraffin-embedded sections may require prolonged proteolytic unmasking (e.g., by pepsin but not by trypsin). The product reacts with human fibronectin. No reactivity is observed with other species including goat, bovine, dog, rabbit, mouse and chicken.

Fibronectin (FN)⁹⁻¹¹ is an extracellular matrix protein composed of two nearly identical, disulfide-bound polypeptides with typical molecular weights of 220-280 kDa. Cellular fibronectin is structurally and antigenically similar to cold insoluble globulin from plasma and antibodies to either form usually crossreact. Careful analysis of the FN molecule indicates that it contains several functionally and structurally distinct domains that may bind to cell surfaces and to a variety of molecules such as collagen, heparin, gelatin, fibrin and DNA. Human fibronectin is composed of at least five distinct domains which are referred to as Hep-1/Fib-1, Gel, Cell, Hep-2 and Fib-2, depending on their affinity for heparin (Hep), gelatin (Gel), the cell surface (Cell) or fibrin (Fib). These domains are aligned from the amino to the carboxyl terminus in the above order and can be separated from each other by mild proteolytic digestion.5

Because of their multiple interactions, FNs play an important role in diverse biological phenomena, including cell adhesion, cell migration, hemostasis and thrombosis, wound healing and the ability to induce a more normal phenotype in transformed cells. Upon malignant transformation, many cells lose most of their surface-bound FN. Immunohistochemical studies show the presence of cellular fibronectin in sections of human malignant tumors, indicating that at least part of the fibronectin in such tumors is not derived from the plasma but is produced locally.^{4,5,12} Monoclonal antibodies reacting specifically with FN may be used to localize FN in human cell cultures, in tissue sections and in plasma, by immunoblotting and immunohistochemical techniques. Also, they may be used for immunoaffinity purification and immunoprecipitation of cellular fibronectin. These antibodies may be used to specifically modulate the functional activity of FN.

Reagents

Supplied as a solution in 0.01 M PBS, pH 7.4, containing 15 mM Sodium azide as a preservative.

Precautions and Disclaimer

Due to the Sodium azide content a safety data sheet (SDS) for this product has been sent to the attention of the safety officer of your institution. Consult the SDS for information regarding hazardous and safe handling practices.

Storage/Stability

1

Store at 2-8 °C for up to one month. For extended storage, freeze in working aliquots. Repeated freezing and thawing is not recommended. Storage in "frost-free" freezers is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use.



Product Profile

A minimum working dilution of 1:100 was determined by indirect immunofluorescent labeling of cultured human fibroblasts. each user determine the optimal working dilution for individual applications by titration assay.

References

- 1. Sekiguchi, K., et al., Biochem. Biophys. Res. Commun., 116, 534 (1983).
- 2. Zardi L., et al., Int. J. Cancer, 25, 325 (1980).
- 3. Zardi, L., et al., EMBO J., 6, 2337 (1987).
- Carnemolla, B., et al., J. Cell Biol., 108, 1139 (1989).
- 5. Zardi, L., Eur. J. Biochem., 146, 571 (1985).
- 6. Borsi, L., et al., J. Cell Biol., 104, 595 (1987).
- 7. Sekiguchi, K., et al., J. Biol. Chem., 260, 5105 (1985).
- 8. Carnemolla, B., et al., In: Prot. Biol. Fluids, H. Peeters (ed.), Pergamon Press, Oxford, 30, 409 (1982).
- Mosher, D. F., Prog. Hemost. Thromb., 5, 111 (1980).
- Yamada, K. M., Annu. Rev. Biochem., 52, 761 (1983).
- 11. Hynes, R., Annu. Rev. Cell Biol., 1, 67 (1985).
- Castellani, P., et al., J. Cell Biol., 103, 1671 (1986).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Terms and Conditions of Sale

Warranty, use restrictions, and other conditions of sale may be found at <u>SigmaAldrich.com/terms</u>.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.



