

Data Sheet

3dGRO™ L-WRN Conditioned Media Supplement

Organoid Culture Medium

SCM105**Pack Size 50 mL****Store at -20 °C****FOR RESEARCH USE ONLY****Not for use in diagnostic procedures. Not for Human or Animal Consumption.**

Background

The 3dGRO™ L-WRN Conditioned Media Supplement is a formulation based on DMEM/F-12 PLUS medium containing the 3 growth factors Wnt3a (W), R-spondin-3 (R), and Noggin (N). These factors act upon the Wnt signaling pathway, which plays a critical role in sustaining the proliferation of LGR5+ crypt stem cells within organoids. Wnt activation is thus essential for in vitro growth of human gut organoids.¹ L-WRN conditioned medium is widely used as a component of the culture medium for supporting the growth of human intestinal organoids² including the 3dGRO™ gastrointestinal organoid biobank (Cat. No. SCC310-SCC367). Applications for L-WRN conditioned medium include but are not limited to the expansion of human intestinal gut organoids and other epithelial organoid cell types.

Source

The 3dGRO™ L-WRN Conditioned Media Supplement is derived from the Mus musculus L-M(TK-) cell line secreting mouse Wnt3a, R-spondin-3 and Noggin. L-WRN Conditioned Medium may contain trace amounts of mouse DNA.

Quality Control Testing

- Appearance (color): clear, red liquid
- Endotoxin: < 2 EU/mL
- Sterility Tested: No Growth/Pass
- Media are tested for function in supporting growth of human intestinal organoids
- Media are tested for levels of growth factors by ELISA

Storage and Handling

The 3dGRO™ L-WRN Conditioned Media Supplement should be stored at -20 °C. When ready to use, thaw the L-WRN Conditioned Medium overnight at 2-8 °C in the dark. Thawed medium is good for two weeks when stored at 2-8 °C. If media will not be used within two weeks, aliquot unused media and store at -20 °C. After thawing refrozen aliquots, do not re-freeze media.

Representative Data




Figure 1. Human colonic organoids grown for 15 passages in complete medium containing 50% L-WRN Conditioned Medium (A). Human duodenal organoids grown for 10 passages in Complete Medium containing 50% L-WRN Conditioned Medium (B).

Protocols

Thawing Medium

1. Thaw medium at 2-8 °C overnight, or at room temperature. To minimize loss of growth factor activity, do not thaw medium at 37 °C.
2. Once thawed, use immediately or aliquot and store at -20 °C.

Preparing 1X Complete Medium for Human Intestinal Organoids

Below is the optimized formulation for culture and expansion of human intestinal organoids. For non-intestinal organoids that are derived from epithelial tissues, L-WRN Conditioned Medium may be used but the 2X Media Supplement for the specific organoid system will have to be optimized by the user.

1. To prepare 100 mL of 1X Complete Medium, 50 mL of L-WRN Conditioned Medium is combined with 50 mL 2X Intestinal Media Supplement. The formulation for 2X Intestinal Media Supplement is below.

2X Intestinal Media Supplement (Total volume = 50 mL)

All items may be purchased at [SigmaAldrich.com](https://www.sigmaaldrich.com) unless otherwise noted.

Component	Volume	2X Conc	Cat. No.
DMEM/F12 Plus Basal Medium	42.6 mL	2X	SCM162
N-2 Supplement (100X)	1 mL	2X	ThermoFisher 17502048
B-27™ Supplement (50X), without vitamin A	2 mL	2X	ThermoFisher 12587010
L-Glutamine Solution (100X), 200 mM	1 mL	2X	TMS-002-C
HEPES Solution, 1M in water	1 mL	20 mM	H0887
Niacinamide, prepared as 1M solution in water	1 mL	20 mM	N0636
N-Acetyl-L-cysteine, prepared as 500 mM solution in water	200 µL	2 mM	A9165
Human EGF, reconstituted to 200 µg/mL in PBS/ 0.1% BSA	50 µL	200 ng/mL	E9644
[Leu ¹⁵]-Gastrin I, reconstituted to 100 µM in PBS/ 0.1% BSA	10 µL	20 nM	G9145
Prostaglandin E2, reconstituted to 100 µM in DMSO	10 µL	20 nM	P6532
A-83-01, reconstituted to 1 mM in DMSO	50 µL	1 µM	SML0788
SB202190, reconstituted to 20 mM in DMSO	50 µL	20 µM	S7067
Penicillin-Streptomycin, 100x solution	1 mL	2X	P4333

2. Filter the 2X Intestinal Media Supplement using a 0.2 µm filter.
3. Combine 50 mL of 2X Intestinal Media Supplement with 50 mL of L-WRN Conditioned Medium; mix well. Use 1X Complete Media within 2 weeks and store at 2-8 °C.
4. For increased viability of human gut organoids during passaging, calculate the total volume of 1X Complete Medium required for passaging and add 2.5 µM CHIR99021 (Cat. No. SML1046) to just that volume. CHIR99021 should only be present for the first two days of each passage. It is not necessary to add CHIR99021 to the 1X Complete Medium during regular media exchanges.
5. Y-27632 (ROCK inhibitor) (Cat. No. SCM075) at 10 µM final concentration is recommended to be added to 1x Complete Medium upon passaging.
6. Primocin® (Invivogen ant-pm-1) at 100 µg/mL final concentration may be added to 1x Complete Medium as an additional antimicrobial supplement.

References

1. *Cell Signal* 2014; 26(3): 570-9.
2. *Gut* 2015; 64(6): 911-920.

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Terms and Conditions of Sale

Warranty, use restrictions, and other conditions of sale may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck KGaA, Darmstadt, Germany, operates as MilliporeSigma in the U.S. and Canada.

MilliporeSigma, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.
© 2023-2024 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

Document Template 20306518 Ver 6.0
20508389 Ver 3.0, Rev 31May2024, CJ

**Millipore
Sigma**