Quick Start Guide

GenElute™ Direct mRNA Kit

DMN10, DMN70

Reagents to Prepare

- Dissolve the Proteinase K in 40% glycerol. Use 0.5 mL of 40% glycerol for 5 mg Proteinase K (10 preps) or 2.5 mL of 40% glycerol for 25 mg Proteinase K (70 preps). Mix thoroughly before use. Store at 2-8 °C.
- Prepare fresh Lysis Solution with Proteinase K by adding 20 µl of Proteinase K solution per mL of Lysis Solution and mixing thoroughly. The required amount per preparation is 0.5 mL for cultured cells and 1.0 mL for tissue.

Protocol

All spins at \geq 12,000 x g.

Release RNA from Cells

- 1. Cultured Cells
 - a. Pellet up to 10⁷ cells. Discard medium.
 - Add 0.5 ml Lysis Solution containing Proteinase K to cell pellet & vortex.
 - c. Transfer lysate to filtration column and spin 2 min.
 - d. Continue to Digest Proteins and Bind mRNA to Beads

Release RNA from Tissues

- 1. Mammalian Tissues
 - a. Place up to 50 mg tissue in homogenizer vessel.
 - b. Add 1 ml Lysis Solution containing Proteinase K and homogenize immediately.
 - c. Continue to Digest Proteins and Bind mRNA to Beads.

Digest Proteins and Bind mRNA to Beads

- 2. Incubate lysate at 65 °C for 10 min.
- 3. Add 32 μ l 5M NaCl to cells OR 64 μ l 5M NaCl to tissue.
- 4. Add 25 μl oligo (dT) beads and vortex.
- 5. Incubate at room temperature for 10 min.
- 6. Spin 5 min to pellet beads. Discard all but ${\sim}50~\mu l$ supernatant.

Optional: Release mRNA & Rebind

- Resuspend beads in 500 μl Lysis Solution + 32 μl 5M NaCl Pulse vortex.
- 8. Incubate at 65 °C for 5 min.
- 9. Incubate at room temperature for 5 min.
- 10. Spin for 2 min to pellet beads. Discard all but $\sim 50~\mu l$ supernatant.

Wash to Remove Contaminants

- 11. Thoroughly resuspend beads in 350 μ l Wash Solution and transfer to spin column.
- 12. Spin 1-2 min. Discard flow-through.
- 13. **2x:** Add 350 μl Low Salt Wash Solution. Spin 1-2 min. Discard flow-through.

Elute Purified mRNA

- 14. Transfer column to new collection tube.
- 15. Add 50 µl Elution Buffer (pre-warmed to 65°C).
- 16. Incubate at 65°C for 2-5 min. Spin 1 min. **Optional:** Repeat elution for additional yield.

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page on our web site at SigmaAldrich.com/TechService.

Terms and Conditions of Sale

Warranty, use restrictions, and other conditions of sale may be found at SigmaAldrich.com/Terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/Offices.

Merck, Millipore, GenElute and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2022 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the U.S. and Canada.

