

Product Information

Anti-MAP2 (2a+2b) Antibody, Mouse Monoclonal

~2 mg/mL, Clone AP-20, purified from hybridoma cell culture

M2320

Product Description

Anti-MAP2 (2a + 2b) (mouse IgG1 isotype) is derived from the hybridoma produced by the fusion of mouse myeloma cells and splenocytes from a mouse immunized with bovine microtubule associated protein 2 (MAP2). The isotype is determined using a double diffusion immunoassay using Mouse Monoclonal Antibody Isotyping Reagents (Cat. No. ISO2).

Anti-MAP2 specificity was established by an immunoblot.¹ The antibody localizes the high molecular weight forms of MAP2, namely MAP2a and MAP2b, but shows no reactivity with MAP2c. No cross-reactivity is observed with MAP1, MAP5, tubulin, or tau (T). The antibody reacts with human,² bovine, rat,^{3,4} mouse,⁵ *Xenopus*, salamander, and quail tissue or cells by immunocytochemical⁴ techniques utilizing either a fluorescent or peroxidase label, or by immunohistochemistry.^{2,3,5}

MAP2 is the major microtubule associated protein of brain tissue. There are three forms of MAP2; two are similarly sized with apparent molecular masses of 280 kDa (MAP2a and MAP2b) and the third with a lower molecular mass of 70 kDa (MAP2c). In the newborn rat brain, MAP2b and MAP2c are present, while MAP2a is absent. Between postnatal days 10 and 20, MAP2a appears. At the same time, the level of MAP2c drops by 10-fold.⁶⁻¹³ This change happens during the period when dendrite growth is completed and when neurons have reached their mature morphology. MAP2 is degraded by a Cathepsin D-like protease in the brain of aged rats. There is some indication that MAP2 is expressed at higher levels in some types of neurons than in other types. MAP2 is known to promote microtubule assembly and to form side arms on microtubules. It also interacts with neurofilaments, actin, and other elements of the cytoskeleton.⁶⁻¹³

Reagent

Supplied as a solution in 0.01 M phosphate buffered saline, pH 7.4, containing 15 mM sodium azide.

Antibody Concentration: ~2 mg/mL

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For extended storage, freeze in working aliquots. Repeated freezing and thawing is not recommended. Storage in "frost-free" freezers is also not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilution samples should be discarded if not used within 12 hours.

Product Profile

Immunoblotting:

A working antibody concentration of 1-3 µg/mL is recommended using a rat brain preparation or rat cerebral cortex extract.

Note: In order to obtain the best results using various techniques and preparations, we recommend determining optimal working dilutions by titration.

References

1. Belanger, D., et al., J. Cell Sci., **115**, 1523-1539 (2002).
2. Almer, G., et al., Brain, **122**, 5-16 (1999).
3. Bury, S.D., and Jones, T.A., J. Neuro., **22**, 8597-8606 (2002).
4. Jolimay, N., et al., J. Neuro. **20**, 9111-9118 (2000).
5. Hevner, R.F., et al., Neuron, **29**, 353-366 (2001).
6. Binder, L.I., et al., Ann. NY Acad. Sci., **466**, 145-166 (1986).
7. Garner, C.C., and Matus, A.J., J. Cell Biol., **106**, 779-783 (1983).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Standard Warranty

The applicable warranty for the products listed in this publication may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the U.S. and Canada.

MilliporeSigma, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.
© 2022 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

M2320pis Rev04/22

**MILLIPORE
SIGMA**