

3050 Spruce Street, St. Louis, MO 63103 USA
Tel: (800) 521-8956 (314) 771-5765 Fax: (800) 325-5052 (314) 771-5757
email: techservice@sial.com sigma-aldrich.com

Product Information

SILu™Prot CST3, Cystatin C, human recombinant, expressed in *E. coli* MS Protein Standard, ¹⁵N-labeled

Catalog Number **MSST0059** Storage Temperature –20 °C

Synonyms: Cystatin 3, CYTC, Post-gamma-globulin, Neuroendocrine basic polypeptide.

Product Description

SILu[™]Prot CST3 is a recombinant, ¹⁵N stable isotopelabeled, human CST3, expressed in *E. coli*. It is a protein of 120 amino acids, with a calculated molecular mass of 13.4 kDa. SILu[™]Prot CST3 is designed to be used as an internal standard for bioanalysis of CST3 in mass spectrometry.

Cystatin C (CST3) is a small protein consisting of a single polypeptide chain, secreted at a constant rate by most nucleated cells in human body. It belongs to the family of papain-like cysteine protease inhibitors, and its main biological role is the inhibition of extracellular cathepsins. Since Cystatin C is constantly secreted, freely filtered by the glomerulus and then fully absorbed in the proximal tubules, the serum concentration is considered to be an accurate marker of the glomerular filtration rate. Cystatin C is normally absent in the urine, therefore it has been suggested that the presence of urinary Cystatin C reflects the level of tubular epithelial damage. Since Cystatin C reflects the level of tubular epithelial damage.

Patients with chronic kidney disease are at high risk for developing cardiovascular disease and cardiovascular events. In recent years Cystatin C has emerged as a potential marker for cardiovascular risk. 4,5

Cystatin C has also been proposed to have a role in modification of the proteolytic system in cancer. Elevated serum cystatin C levels in cancer patents, are linked to meager outcome of disease. ⁶

Each vial contains 10 μ g of SILuTMProt CST3 standard, lyophilized from a Tris buffered solution, pH 8.0, containing 10 mM methionine as stabilizer. Vial content is determined according to protein absorbance at 280 nm (ϵ = 0.877).

Purity: ≥95% (SDS-PAGE)

Heavy nitrogen incorporation efficiency: ≥97% (MS)

UniProt: P01034

Sequence Information:

SSPGKPPRLVGGPMDASVEEEGVRRALDFAVGEYN KASNDMYHSRALQVVRARKQIVAGVNYFLDVELGRT TCTKTQPNLDNCPFHDQPHLKRKAFCSFQIYAVPWQ GTMTLSKSTCQDA

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Preparation Instructions

Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein in sterile ultrapure water to a final concentration of 100 μ g/mL.

Storage/Stability

Store the lyophilized product at -20 °C. The product is stable for at least 2 years as supplied.

After reconstitution, it is recommended to store the protein in working aliquots at –20 °C.

References

- Kolodziejczyk, R. et al., Crystal structure of human cystatin c stabilized against amyloid formation. FEBS, 277(7), 1726-1737 (2010).
- 2. Åkerblom, A., Biomarkers of renal function in Acute Coronary Syndromes. *Digital Comprehensive Summaries of Uppsala Dissertations from the Faculty of Medicine* (2013).
- 3. Jiang, R. et al., Detection of Cystatin C biomarker for a clinical measurement of renal disease by developed ELISA diagnostic kits. *J.Trans.Med.*,**12**, 1-8 (2014).
- 4. Angelidis, C. et al., Cystatin–C: an emerging biomarker in cardiovascular diseases. *Curr. Top. Med. Chem.*, 13(2), 164-179 (2013).
- 5. Taglieri, N. et al., Cystatin C and the cardiovascular risk. *Clin. Chem.*, **55(11)**, 1932-1943 (2009).
- 6. Hammouda, N.E. et al., Serum Cystatin C as biomarker in Diffuse Large b-cell Lymphoma. *Sci. Pharm.*, **85(1)**, 1-10 (2017).

Legal Information

Sold under license from DuPont, U.S. Patent No. 7,396,688.

This product is licensed under U.S. Patent No. 7,396,688 and foreign counterparts from E. I. du Pont de Nemours and Company. The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product for research and development only, including services for a third party for consideration. The buyer cannot sell or otherwise transfer this product, its components or materials made using this product or its components to a third party. Information about licenses for excluded uses is available from: E. I. du Pont de Nemours and Company; Attn: Associate Director, Commercial Development; DuPont Experimental Station E268; 200 Powdermill Rd.; Wilmington, DE 19803; 1-877-881-9787 (voice), 1-302-695-1437 (fax), licensing@dupont.com.

SILu is a trademark of Sigma-Aldrich Co. LLC.

DS,JK,MAM 08/17-1