

## RAT ANTI-HUMAN PRO-COLLAGEN I AMINO-TERMINAL MONOCLONAL ANTIBODY

CATALOG NUMBER: MAB1912

LOT NUMBER:

**QUANTITY**:  $100 \mu$ L

**SPECIFICITY:** Human pro-collagen I. MAB1912 labels the amino-terminal pro-peptide.

**IMMUNOGEN:** Human pro-collagen I.

**ISOTYPE**: IgG<sub>1</sub>

CELL LINE: M-58

**APPLICATIONS:** Immunohistochemistry: 1:10,000-1:50,000 (fresh frozen tissue sections.)

Suitable for formaldehyde/formalin fixed paraffin sections, suggested starting dilution 1:100-1:1000, after treatment with 1% trypsin, 20 minutes at room temperature. Does not stain mature collagen fibers in tissue. Localized intracellularly in cells producing pro-collagen I.

Not recommended for Western Blots

Optimal working dilutions must be determined by end user.

FORMAT: Tissue Culture Supernatant

PRESENTATION: Liquid

**STORAGE/HANDLING:** Maintain frozen at -20°C in undiluted aliquots for up to 12 months.

**REFERENCES:** Diease Management and Clinical Outcomes (1998). **1(4):** 136-141.

Important Note: During shipment, small volumes of product will occasionally become entrapped in the seal of the product vial. For

products with volumes of 200 uL or less, we recommend gently tapping the vial on a hard surface or briefly

centrifuging the vial in a tabletop centrifuge to dislodge any liquid in the container's cap.

FOR RESEARCH USE ONLY; NOT FOR USE IN DIAGNOSTIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION

Unless otherwise stated in our catalog or other company documentation accompanying the product(s), our products are intended for research use only and are not to be used for any other purpose, which includes but is not limited to, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses or any type of consumption or application to humans or animals.

©2002 - 2010: Millipore Corporation. All rights reserved. No part of these works may be reproduced in any form without permission in writing.