

## Product Information

## Minimum Essential Medium Eagle (MEM)

## Spinner Modification (S-MEM)

## M8167

Minimum Essential Medium (MEM), developed by Harry Eagle, is one of the most widely used of all synthetic cell culture media. Early attempts to cultivate normal mammalian fibroblasts and certain subtypes of HeLa cells revealed they had specific nutritional requirements that could not be met by Eagle's Basal Medium (BME). Subsequent studies using these and other cells in culture indicated additions to BME could be made to aid growth of a wider variety of fastidious cells.

MEM, which incorporates these modifications, includes higher concentrations of amino acids so the medium more closely approximates the protein composition of cultured mammalian cells. MEM has been used for cultivation of a wide variety of cells grown in monolayers. Optional supplementation of non-essential amino acids to the formulations that incorporate either Hanks' or Earle's salts has broadened the usefulness of this medium. The formulation has been further modified by optional elimination of calcium to permit growth of cells in suspension culture.

| COMPONENT                                    | M8167<br>[1x] g/L |
|----------------------------------------------|-------------------|
| <b>Inorganic Salts</b>                       |                   |
| MgSO <sub>4</sub> (anhydrous)                | 0.09767           |
| KCl                                          | 0.4               |
| NaCl                                         | 6.8               |
| Na <sub>2</sub> HPO <sub>4</sub> (anhydrous) | 1.22              |
| <b>Amino acids</b>                           |                   |
| L-Arginine • HCl                             | 0.126             |
| L-Cystine • 2HCl                             | 0.0313            |
| L-Histidine • HCl • H <sub>2</sub> O         | 0.042             |
| L-Isoleucine                                 | 0.052             |
| L-Leucine                                    | 0.052             |
| L-Lysine • HCl                               | 0.0725            |
| L-Methionine                                 | 0.015             |
| L-Phenylalanine                              | 0.032             |
| L-Threonine                                  | 0.048             |
| L-Tryptophan                                 | 0.01              |
| L-Tyrosine • 2Na • 2H <sub>2</sub> O         | 0.0519            |
| L-Valine                                     | 0.046             |
| <b>Vitamins</b>                              |                   |
| Choline Chloride                             | 0.001             |
| Folic Acid                                   | 0.001             |
| <i>myo</i> -Inositol                         | 0.002             |
| Niacinamide                                  | 0.001             |
| D-Panthenic Acid • ½Ca                       | 0.001             |
| Pyridoxal • HCl                              | 0.001             |
| Riboflavin                                   | 0.0001            |
| Thiamine • HCl                               | 0.001             |

| <b>COMPONENT</b>   | <b>M8167</b><br>[1x] g/L |
|--------------------|--------------------------|
| <b>Other</b>       |                          |
| Glucose            | 1                        |
| Phenol Red • Na    | 0.011                    |
| NaHCO <sub>3</sub> | 2.2                      |
| <b>ADD</b>         |                          |
| L-Glutamine        | 0.292                    |

## References

1. Eagle, H., et al., myo-Inositol as an Essential Growth Factor for Normal and Malignant Human Cells in Tissue Culture. J. Biol. Chem., 214, 845- 847(1956).
2. Eagle, H., Media for Animal Cell Culture. Tissue Culture Association Manual, 3, 517-520 (1976).
3. Eagle, H., Amino Acid Metabolism in Mammalian Cell Cultures. Science, 130, 432-437 (1959).
4. Eagle, H., Nutrition Needs of Mammalian Cells in Culture. Science, 122, 501 (1955).

## Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

### Technical Assistance

Visit the tech service page at [SigmaAldrich.com/techservice](https://SigmaAldrich.com/techservice).

### Terms and Conditions of Sale

Warranty, use restrictions, and other conditions of sale may be found at [SigmaAldrich.com/terms](https://SigmaAldrich.com/terms).

### Contact Information

For the location of the office nearest you, go to [SigmaAldrich.com/offices](https://SigmaAldrich.com/offices).

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the U.S. and Canada.

MilliporeSigma, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2024 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.  
M8167pis Rev 11/24

**Millipore**  
**SIGMA**