

# **Technical Data Sheet**

# Egg-Yolk Tellurite Emulsion sterile, 20 %, for microbiology

Ordering number: 1.03785.0001

Egg-yolk tellurite emulsion is used as an additive, e.g. for GranuCult™ BAIRD-PARKER agar (base) acc. ISO 6888 and FDA-BAM (article number 1.05406.0500).

#### **Mode of Action**

Egg-yolk permits the detection of microbial lecithinase activity and tellurite reduction.

## **Typical Composition**

Sterile Egg-Yolk	200 ml/l
NaCl	4.25 g/l
Potassium tellurite	2.1 g/l
distilled water to give a final volume of 1000 ml	

#### **Preparation**

Shake the bottle well to suspend any sediment. Mix 50 ml with 950 ml of the reconstituted culture. **Caution:** Observe sterile procedure when emptying the bottle

#### **Storage**

Usable up to the expiry date when stored dry and tightly closed at +2 °C to +8 °C.

## **Quality Control**

Egg-Yolk Emulsion is tested in GranuCult™ BAIRD-PARKER agar (base) acc. ISO 6888 and FDA-BAM (article number 1.05406.0500)

Function	Control strains	Incubation	Reference medium	Method of control	Expected results
Productivity	<i>Staphylococcu</i> <i>s aureus</i> ATCC® 6538	22-26 h to - 46-50 h at 36-38 °C		Quantitative	Recovery ≥ 50 %, black
	<i>Staphylococcu</i> <i>s aureus</i> ATCC® 25923				or grey colonies with clear halo



1.03785.0001 Page 1 of 3

Function	Control strains	Incubation	Reference medium	Method of control	Expected results
Selectivity	Escherichia coli ATCC® 8739	- 46-50 h at 36-38 °C	-	Qualitative	No recovery limit, black or
	Escherichia coli ATCC® 25922				grey colonies without clear halo
Specificity	Staphylococcus saprophyticus ATCC® 15305	- 22-26 h to 46-50 h at - 36-38 °C	t -	Qualitative	No recovery limit, brown
	Staphylococcu s epidermidis ATCC® 12228				or black colonies without clear
	Enterococcus hirae ATCC® 8043				halo
	Proteus mirabilis ATCC® 29906				No recovery limit, brown or black colonies without clear halo

Please refer to the actual batch related Certificate of Analysis.

The performance test is in accordance with the current version of EN ISO 11133.

A recovery rate of 50 % is equivalent to a productivity value of 0.5.

#### Literature

APHA (2015): Compendium of Methods for the Microbiological Examination of Foods.  $5^{th}$  ed. American Public Health Association, Washington, D.C.

FDA-BAM (2001): Chapter No. 12: *Staphylococcus aureus*. U.S. Food and Drug Administration - Bacteriological Analytical Manual.

Baird-Parker, A.C. (1962): An improved diagnostic and selective medium for isolating coagulase positive staphylococci. J. Appl. Bact. **25**: 12-19.

ISO International Standardisation Organisation. Microbiology of food and animal feeding stuffs -- Horizontal method for the enumeration of coagulase-positive staphylococci (*Staphylococcus aureus* and other species) - Part 1: Technique using Baird-Parker agar medium + Amendment 1: Inclusion of precision data. EN ISO 6888-1:1999/Amd 1:2003.

ISO International Standardisation Organisation. Microbiology of food and animal feeding stuffs -- Horizontal method for the enumeration of coagulase-positive staphylococci (*Staphylococcus aureus* and other species) - Part 3: Detection and MPN technique for low numbers. EN ISO 6888-3:2003.

ISO International Standardisation Organisation. Microbiology of food, animal feed and water - Preparation, production, storage and performance testing of culture media. EN ISO 11133:2014.



1.03785.0001 Page 2 of 3

## **Ordering Information**

Product	Cat. No.	Pack size
Egg-Yolk Tellurite Emulsion 20% (sterile)	1.03785.0001	10 x 50 ml
GranuCult™BAIRD-PARKER Agar (Base) acc. ISO 6888 and FDA-BAM	1.05406.0500	500 g

MilliporeSigma 290 Concord Road Billerica, MA 01821 Find contact information for your country at: www.EMDmillipore.com/offices

For Technical Service, please visit: www.EMDmillipore.com/techservice

For more information, visit

www.EMDmillipore.com/biomonitoring







1.03785.0001 Page 3 of 3