

Product Information

CRYO-DMSO-F

Liquid, Sterile-Filtered, 100 mL

C9252

Introduction

CRYO-DMSO-F is a serum-free, protein free, animal origin-free, DMSO Free and fully defined cryopreservation medium. Designed to prepare and preserve cells in ultralow temperature environments (–80 to –196 °C), It provides a safe, protective environment for cells during the freezing, storage, and thawing process. CRYO-DMSO-F contained proprietary components which are directly reducing the level of freezing induced apoptosis and necrosis and improving post-thaw cell viability and function. CRYO-DMSO-F (C9252) is recommended for the preservation of Primary cells, iPSC, MS, MSC, Oocyte, Embryo, CAR-T, NK, and Macrophage cells. It is cGMP-mimic condition manufactured with high quality grade components.

Appearance (Turbidity)

Clear liquid

Appearance (Form)

Solution

Sterility

Sterile Filtered (0.22 µM)

pH

7.80~8.20

Preparation instructions

The CRYO-DMSO-F solution is ready-to-use and complete with no additives required. Wipe down the outside of container with 70% alcohol before opening as the contents are sterile. If the seal has been broken, do not use it.

Storage/Stability

Store the CRYO-DMSO-F solution at 2-8 °C and protected from light until ready to use it.

Protocol

Freezing Procedures

1. Suspended cell to be cryopreserved using mechanically or enzymatically dissociation.
2. Centrifuge the cells to obtained pellet.
3. Remove supernatant (remove the culture media as possible to reduce dilution of the CRYO-DMSO-F solution).
4. Add Ambient temperature CRYO-DMSO-F solution to a cell concentration range of $1-10 \times 10^7$ cells /1 mL of CRYO-DMSO-F for standard cell culture protocol.
5. After mixed with CRYO-DMSO-F solution with cells, incubated for 10 minutes at 1-4 °C to penetrate cryoprotectants inside of cells (in case of small tissue or organoid, 20 minutes incubation at 1-4 °C).

6. Nucleation-lower sample temperature -80°C ; After cells are mixed with solution, put cryovial into controlled rate freezer ($-1^{\circ}\text{C}/\text{minute}$) and then freeze to -80°C (slow freezing method), or put cryovial into Bicell, Mr. Frosty[®] Freezing container, or similar kinds of slow freezing container and put the cryovial included in such slow freezing container into -80°C freezer.

NOTE: After finished the nucleation of cells at -80°C freezer or controlled rate freezer, store the freeze cell at liquid nitrogen tank (below -130°C).

Thawing Procedures

1. Thaw samples quickly in a 37°C water bath. Samples should be thawed with gentle swirling of the sample until all visible ice has melted (Do not allow sample to warm above chilled temperatures ($0-10^{\circ}\text{C}$). Cryovials should be cool to the touch when removed from the water bath).
2. Dilute cell/CRYO-DMSO-F mixture immediately with appropriate culture medium. Add 20 mL of culture medium to 50 mL conical tube and gently mix with thawed cell/CRYO-DMSO-F mixture to culture medium (The dilution culture medium should be $20-37^{\circ}\text{C}$). A dilution ratio of 1:20 V:V (1 mL of thaw cell + 20 mL of culture media) or greater is recommended. After diluting cells with warmed culture medium, gently inverted for 5-10 times and follow by centrifuge (recommend at $400 \times g$ for 5 minutes).
3. After centrifuge, completely suction of supernatant and add new warmed culture media.
4. Plate the cells appropriately and cultured the cells or use immediately.

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Terms and Conditions of Sale

Warranty, use restrictions, and other conditions of sale may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck operates
as MilliporeSigma in the U.S. and Canada.

Merck, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2025 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.
23235063# Rev 07/25

