

Product Information

Anti-Monkey IgG (Whole Molecule)-Peroxidase

Antibody Produced in Rabbit, Buffered Aqueous Solution

A2054

Product Description

Antiserum is produced in rabbit using purified monkey IgG (rhesus) as the immunogen. Antibody is isolated from rabbit anti-monkey IgG antiserum by immunospecific purification, which removes essentially all rabbit serum proteins, including immunoglobulins, which do not specifically bind to monkey IgG. Rabbit anti-monkey IgG is conjugated to horseradish peroxidase by protein crosslinking with glutaraldehyde.

Identity and purity of the antibody is established by immunoelectrophoresis (IEP), prior to conjugation. Electrophoresis of the antibody preparation followed by diffusion versus anti-rabbit IgG and anti-rabbit whole serum result in single arcs of precipitation.

Reagent

Provided as a solution in 0.01 M phosphate buffered saline, pH 7.4, containing 0.05% MIT as a preservative.

Antibody concentration: 4-11 mg/mL

Molar Ratio: (Antibody:Peroxidase): 0.6-1.5

Precautions and Disclaimer

For research use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For extended storage, the solution may be frozen in working aliquots. Repeated freezing and thawing, or storage in "frost-free" freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use.

Product Profile

Direct ELISA

A working dilution of at least 1:45,000 is determined using 5 µg/mL monkey IgG for the coat and OPD, Cat. No. P8287, as substrate.

Dot Blot

A dilution of 1:160,000 was determined in a direct chemiluminescence assay using 10 ng monkey IgG/dot. Luminol plus enhancer was used as substrate.

Note: Working dilutions should be determined by titration assay. Due to differences in assay systems, these titers may not reflect the user's actual working dilution.

References

1. Voller, A. et al., *Bulletin WHO*, **53**: 55 (1976).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Standard Warranty

The applicable warranty for the products listed in this publication may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.

Merck and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.
© 2022 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.