Sigma-Aldrich_®

Product Information

Apyrase from potatoes

ATPase ≥200 units/mg protein, lyophilized powder

A6535

Product Description

CAS Number: 9000-95-7

Enzyme Commission (EC) Number: 3.6.1.5

Synonyms: Adenosine 5'-diphosphatase, Adenosine 5'-triphosphatase, ATP diphosphohydrolase¹

Apyrase has adenosine 5'-triphosphatase (ATPase) and adenosine 5'-diphosphatase (ADPase) activities,² where hydrolysis of the pyrophosphate bonds leads to sequential release of inorganic orthophosphate.³ At least two isoenzymes with different ATPase/ADPase ratios exist in different varieties of potato (*Solanum tuberosum*):^{4,5}

- 'Pimpernel' isoenzyme
 - High ATPase/ADPase ratio of ~10:1
 - o Isoelectric point (pI) of 8.74
- 'Desirée' isoenzyme
 - Low ATPase/ADPase ratio of ~1:1
 - Isoelectric point (pI) of 6.69

These isozymes each have a molecular mass of ~49 kDa (gel filtration).⁵ This product is predominantly the low ATPase/ADPase ratio isozyme.

Apyrase requires divalent metal ions for activity, with optimal activity observed with 5 mM Ca²⁺.

- For hydrolysis of organic diphosphates and triphosphates, the optimal pH is 6.5
- For inorganic substrates, the optimal pH is 5.1.⁵

Several theses^{6,7} and dissertations⁸⁻¹⁵ have cited use of product A6535 in their research protocols.

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Preparation Instructions

This product is soluble in water (1 mg/mL). One publication reports preparation of 125 units/mL stock solutions of this product in 10 mM HEPES (pH 7.8).¹⁶

Storage/Stability

- Stock solutions at pH between 5-7.5 can be stored as frozen aliquots, such as in 30 mM HEPES, pH 7.2.¹⁷
- For enzyme solutions of <1 mg/mL, dissolve in HEPES buffer (pH 7.5) containing 1 mM MgCl₂, 1 mM DTT, 1 mM EDTA, and 1 mg/mL BSA.
- Repeated freeze-thaw cycles risk loss of activity.
- One publication indicates storage of 340 units/mL stock solutions of apyrase in calcium-free Tyrode's buffer, at -20 °C, in single-use aliquots.¹⁸

References

1

- Vasconcelos, E.G. et al., "Potato apyrase: a new tool for parasitic disease research", in Potato II. Fruit, Vegetable and Cereal Science and Biotechnology 3 - Special Issue 1 (P. Tennant and N. Benkeblia, eds.). Global Science Books, pp. 80-88 (2009).
- 2. Traverso-Cori, A. et al., Arch. Biochem. Biophys., **109(1)**, 173-184 (1965).



- Mancilla, M. et al., Phytochemistry, 23(7), 1397-1400 (1984).
- 4. Molnar, J., and Lorand, L., *Arch. Biochem. Biophys.*, **93(2)**, 353-363 (1961).
- Kettlun, A. et al., Phytochemistry, 21(3), 551-558 (1982).
- 6. Clarke, Andrea, "Externalized ATP in intact erythrocyte suspensions: modulation and determination". Concordia University, M.Sc. thesis, pp. 50, 54 (2012).
- 7. de San Eustaquio Campillo, Alba, "Studies on the expression of recombinant resistance proteins domains". University of Durham, M.Sc. thesis, p. 31 (2012).
- Haserück, Nadine, "Platelet activation and platelet-monocyte aggregate formation induced by the atherosclerotic plaque lipid lysophosphatidic acid". Ludwig-Maximilians-Universität München, Dr. med. dissertation, p. 16 (2007).
- 9. Chen, Taotao, "DnaK Functions as a Central Hub in the *E. coli* Chaperone Network". Ludwig-Maximilians-Universität München, Ph.D. dissertation, pp. 55, 56 (2011).
- 10. Liu, Yiwei, "Non-platelet Purinergic Signaling and Tissue Factor Expression". Auburn University, Ph.D. dissertation, p. 36 (2015).
- McIntosh, Betsy Buechler, "Engineered Cytoskeletal Arrays Reveal Mechanisms of Membrane Transport and Tubulation". University of Pennsylvania, Ph.D. dissertation, p. 114 (2017).
- 12. Smith, Christopher William, "Regulation of platelet production and function by ITIM-containing receptors". University of Birmingham, Ph.D. dissertation, p. 47 (2018).
- 13. Wei, Hao, "Mechanisms of microparticle release from human platelets". University of Cambridge, Ph.D. dissertation, p. 49 (2019).

- 14. Kaschubowski, Klaus Eric, "Use of Fluorescent Sensors to Visualize P2X7-Mediated Changes in Local ATP Concentrations in the Cytosol and at the Cell Surface". Universität Hamburg, Dr. rer. hum. biol. dissertation, p. 25 (2020).
- 15. Dickhout, Annemiek, "Painting proteins to study cardiovascular pathology". Universiteit Maastricht, Ph.D. dissertation, pp. 97, 125 (2022).
- 16. Gonda, K., and Kikyo, N., *Methods Mol. Biol.*, **348**, 247-258 (2006).
- 17. Gennerich, A., and Reck-Peterson, S.L., *Methods Mol. Biol.*, **783**, 63-80 (2011).
- 18. Rutledge, T.W., and Whiteheart, S.W., *Methods Mol. Biol.*, **272**, 109-120 (2004).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Terms and Conditions of Sale

Warranty, use restrictions, and other conditions of sale may be found at <u>SigmaAldrich.com/terms</u>.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.

Merck and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

