

Product Information

ExoPolyzyme

For Enzymatic Lysis of Extracellular Biofilm Matrix

MBD0064

Product Description

Biofilms are complex structures, which adhere to surfaces and comprise of one or more types of bacteria and fungi. They form an extracellular matrix (ECM) which is generally composed of secreted polymers such as exopolysaccharides (EPS), extracellular DNA (eDNA), peptides and lipids.¹

Biofilms may form on a wide variety of biotic and abiotic surfaces including living tissues, medical devices (such as catheters), water system pipes, plants (roots) and aquatic systems.^{2,3}

The EPS produced by certain bacterial strains forms a biofilm that provides resistance to antibiotics and increased tolerance to harsh conditions, such as pH, osmolarity, temperature, nutrients scarcity, mechanical and shear forces.^{4,5}

In addition to being a major issue in hospital and industry settings, the biofilm structure has also been shown to pose a problem in the field of Metagenomics. Many microorganisms present in natural environments reside in biofilms and are therefore not accessible using standard DNA extraction protocols. Mechanical methods used for cell lysis can increase the efficiency of DNA extraction, but can also break the DNA, resulting in lower quality and quantity of DNA extracted from biofilm samples. This may, in turn, introduce bias into the analysis and the results will not represent the whole population of bacteria in the sample.⁶⁻⁸

The ExoPolyzyme is a mixture of seven enzymes that break down different components residing in ECM, such as EPS's and lipids. Each of these enzymes has been previously used to inhibit or remove biofilm.⁹⁻¹⁵

It can be used in biofilm research and as a gentler method for removing the EPS and lipids from the ECM

in metagenomic protocols, providing better access to the bacterial cells within the biofilm and eDNA.

Data has shown that cell lysis using gentle methods, such as MetaPolyzyme (MAC4L) can provide longer DNA molecules that are suitable for sequencing applications.^{16,17}

ExoPolyzyme components:

- α -Amylase
- Cellulase
- β -Glucosidase
- Lyticase
- Alginate lyase
- Lipase
- Hemicellulase

α -Amylase (from *Bacillus* sp.)

α -Amylase belongs to the glycoside hydrolase family 13. It is an enzyme that hydrolyzes α -1,4 glycosidic bonds present in the inner part (endoamylase) of the amylose, or amylopectin chain such as starch and glycogen.¹⁸

Cellulase (from *Trichoderma* sp.)

Cellulases are responsible for cellulose degradation by hydrolyzing the β -1,4-glycosidic bonds. It catalyzes cellulose, hemicellulose, lichenin, and cereal β -D-glucans molecules into monosaccharides such as β -glucose, or shorter polysaccharides and oligosaccharides.¹⁹

β -Glucosidase (from almonds)

β -Glucosidase (also known as β -D-Glucoside glucohydrolase) is a glycoprotein that hydrolyzes terminal, non-reducing β -D-glucosyl residues and releases β -D-glucose.²⁰

Lyticase (from *Arthrobacter luteus*)

Lyticase catalyzes fungal cell wall by β -1,3-glucanase and β -1,3-glucan laminaripentaohydrolase activities.²¹

Alginate lyase

Alginate lyase degrades alginate through elimination of the glycosidic bond. It results in numerous oligosaccharides with unsaturated uronic acid at the non-reducing terminus and unsaturated uronic acid monomers.²²

Lipase (from *Candida rugosa*)

Lipase breaks down triglycerides into free fatty acids and glycerol by catalyzing the hydrolysis of the ester bonds in triglycerides.²³

Hemicellulase (from *Aspergillus niger*):

Hemicellulase is a group of enzymes that includes glucanases, xylanases and mannanase. It's responsible for the degradation of hemicelluloses, which are heteropolysaccharides such as various xylans and mannans by catalyzing β -1,4-glycosidic bonds.²⁴

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

Store this product at -20 °C.

Preparation Instructions

1. Dissolve one vial in 4 ml PBS x1. The PBS should be free of calcium chloride and magnesium chloride (D8537).
2. This amount can be used for approximately 25 reactions in a 96 well plate (150 μ L per well). However, individual optimization should be performed for each biofilm assay.
3. Incubate at 37 °C for at least two hours for optimal activity.

Note: The ExoPolyzyme results in non-specific background in crystal violet biofilm assays. Heat inactivated ExoPolyzyme (95 °C for 10 minutes) may be used as a blank control for crystal violet assays.

References

1. Flemming, Hans-Curt, et al. "Biofilms: an emergent form of bacterial life." *Nature Reviews Microbiology* 14.9 (2016): 563-575.
2. Donlan, Rodney M., and J. William Costerton. "Biofilms: survival mechanisms of clinically relevant microorganisms." *Clinical microbiology reviews* 15.2 (2002): 167-193.
3. Hall-Stoodley, Luanne, J. William Costerton, and Paul Stoodley. "Bacterial biofilms: from the natural environment to infectious diseases." *Nature reviews microbiology* 2.2 (2004): 95-108.
4. Costerton, J. William, Philip S. Stewart, and E. Peter Greenberg. "Bacterial biofilms: a common cause of persistent infections." *science* 284.5418 (1999): 1318-1322.
5. Stewart, Philip S., and J. William Costerton. "Antibiotic resistance of bacteria in biofilms." *The lancet* 358.9276 (2001): 135-138.
6. Moss, Eli L., Dylan G. Maghini, and Ami S. Bhatt. "Complete, closed bacterial genomes from microbiomes using nanopore sequencing." *Nature biotechnology* 38.6 (2020): 701-707.
7. Poptsova, Maria S., et al. "Non-random DNA fragmentation in next-generation sequencing." *Scientific reports* 4.1 (2014): 4532.
8. Knierim, Ellen, et al. "Systematic comparison of three methods for fragmentation of long-range PCR products for next generation sequencing." *PloS one* 6.11 (2011): e28240.
9. Craigen, Bradford, Aliza Dashiff, and Daniel E. Kadouri. "The use of commercially available alpha-amylase compounds to inhibit and remove *Staphylococcus aureus* biofilms." *The open microbiology journal* 5 (2011): 21.
10. Loisele, Melanie, and Kimberly W. Anderson. "The use of cellulase in inhibiting biofilm formation from organisms commonly found on medical implants." *Biofouling* 19.2 (2003): 77-85.
11. Banar, Maryam, et al. "The efficacy of lyticase and β -glucosidase enzymes on biofilm degradation of *Pseudomonas aeruginosa* strains with different gene profiles." *BMC microbiology* 19 (2019): 1-10.
12. Ghadam, Parinaz, Fatemeh Akhlaghi, and Ahya Abdi Ali. "One-step purification and characterization of alginate lyase from a clinical *Pseudomonas aeruginosa* with destructive activity on bacterial biofilm." *Iranian journal of basic medical sciences* 20.5 (2017): 467.
13. Singh, Rahul, et al. "Affordable oral health care: dental biofilm disruption using chloroplast made

- enzymes with chewing gum delivery." *Plant biotechnology journal* 19.10 (2021): 2113-2125. Lee, Jin-Hyung, Yong-Guy Kim, and Jintae Lee. "Thermostable xylanase inhibits and disassembles *Pseudomonas aeruginosa* biofilms." *Biofouling* 34.3 (2018): 346-356.
14. Blakeman, Jamie T., et al. "Extracellular DNA provides structural integrity to a *Micrococcus luteus* biofilm." *Langmuir* 35.19 (2019): 6468-6475.
 15. Tighe, Scott, et al. "Genomic methods and microbiological technologies for profiling novel and extreme environments for the extreme microbiome project (XMP)." *Journal of biomolecular techniques: JBT* 28.1 (2017): 31.
 16. Zhang, Lei, et al. "Comparison analysis of different DNA Extraction methods on suitability for long-read metagenomic nanopore sequencing." *Frontiers in Cellular and Infection Microbiology* 12 (2022): 919903.
 17. Tiwari, S. P., et al. "Amylases: an overview with special reference to alpha amylase." *J Global Biosci* 4.1 (2015): 1886-1901.
 18. Acharya, A., et al. "Isolation and screening of thermophilic cellulolytic bacteria from compost piles." *Scientific world* 10.10 (2012): 43-46.
 19. Singh, Gopal, A. K. Verma, and Vinod Kumar. "Catalytic properties, functional attributes and industrial applications of β -glucosidases." *3 Biotech* 6 (2016): 1-14.
 20. Ribeiro, Ricardo A., et al. "Yeast adaptive response to acetic acid stress involves structural alterations and increased stiffness of the cell wall." *Scientific reports* 11.1 (2021): 12652.
 21. Ertesvåg, Helga, et al. "Biochemical properties and substrate specificities of a recombinantly produced *Azotobacter vinelandii* alginate lyase." *Journal of bacteriology* 180.15 (1998): 3779-3784.
 22. Salah, Riadh Ben, et al. "Biochemical and molecular characterization of a lipase produced by *Rhizopus oryzae*." *FEMS microbiology letters* 260.2 (2006): 241-248.
 23. Koshland Jr, D. E. "Stereochemistry and the mechanism of enzymatic reactions." *Biological reviews* 28.4 (1953): 416-436.

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Terms and Conditions of Sale

Warranty, use restrictions, and other conditions of sale may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.

Merck and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2024 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

MBD0064pis Rev08/24

