

Product Information

Laminin from Engelbreth-Holm-Swarm Murine Sarcoma (Basement Membrane)

L2020

Storage Temperature –20 °C

CAS RN: 114956-81-9

Product Description

Laminin is isolated as the intact native form from Engelbreth-Holm-Swarm tumor, a mouse tumor that produces large amounts of basement membranes.¹ Laminin chains are designated A (molecular weight 400 kDa), B1 (molecular weight 210 kDa) and B2 (molecular weight 200 kDa). The cohesion between these chains is the result of many inter- and intrachain disulfide bonds. Together, they cause the molecule to look like a crucifix. Laminin has active domains for collagen binding, cell adhesion, heparin binding, and neurite outgrowth (PHD) fragment.²

It is recommended for use as a cell culture substratum at 1-2 µg/cm². The optimal concentration will depend upon the cell type as well as the application or research objectives.

Reagent

Supplied at a concentration of approximately 1-2 mg/mL in 50 mM Tris-HCl, pH 7.5, with 150 mM NaCl. It is sterile-filtered using a 0.2 µm filter.

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Preparation Instructions

The solution should be thawed slowly at 2-8 °C. If warmed too quickly, it may form a gel and cannot be reactivated for use.

Storage/Stability

Store unused portion in working aliquots at –20 °C.

Procedure

Optimal conditions for attachment must be determined for each cell line and application.

1. Slowly thaw the laminin solution at 2-8 °C to avoid the formation of a gel.
2. Dilute in a sterile balanced salt solution, or in sterile water, and coat culture surface with a minimal volume.
3. Incubate at 37 °C for two hours
4. Wash 3 times with PBS and plate the cells.

Laminin coatings, such as laminin-coated cover slips, may be stored for approximately one month at 2-8 °C. **Do not use** the product if discoloration or spider web formations appear on the surface of the coated material.

References

5. Timpl, R., et al., Laminin – a glycoprotein from basement membranes. *J. Biol. Chem.*, **254**: 9933- 9937 (1979).
6. Kleinman, H. K., et al., Use of extracellular matrix components for cell culture. *Anal. Biochem.*, **166**: 1- 13 (1987).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Standard Warranty

The applicable warranty for the products listed in this publication may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the U.S. and Canada.

MilliporeSigma, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.
© 2022 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

L2020 Rev 02/22