

Hazard profile of Deviron® 13-S9 detergent for GHS classification


1. Introduction

Deviron® 13-S9 detergent is a fatty alcohol ethoxylate which contains an alkyl chain of 11 to 15 carbon atoms and an ethoxylate chain of 9 ethylene oxide (EO) units. It therefore fulfils the REACH definition of polymer, i.e., *a molecule that contains a sequence of at least 3 monomer units, which are covalently bound to at least one other monomer unit or other reactant* (ECHA, 2023). At the time of writing, polymers are exempted from the provisions of registration under REACH (ECHA, 2023).

The present summary collects the key toxicological and ecotoxicological data that were used for the **GHS hazard classification** of Deviron® 13-S9 detergent. In such evaluation, toxicological and ecotoxicological data available in the public literature on the alkyl PEG ethers group of chemicals were taken into account, based on the documented rationale for grouping fatty alcohol alkoxyates (or alkyl PEG ethers). However, a greater attention was given to alcohol ethoxylates showing an alkyl chain length ranging from C₁₁ to C₁₅, and an EO chain of 9 units. (Q)SAR predictions for several endpoints were also generated for the homologues of Deviron® 13-S9 detergent and considered within a weight of evidence approach.

The weight-of-evidence assessment undertaken for the proposed GHS Classification is described in a more detailed report, which can be provided upon request.

2. Chemical identity

Trade Name	Deviron® 13-S9 detergent
Chemical Name	Alcohols, C11-15-secondary, ethoxylated
Alternative chemical name	C ₁₁₋₁₅ sec-Pareth-9 (C ₁₁₋₁₅ EO ₉)
CAS No.	68131-40-8*
Alternative CAS No./Chemical Name	84133-50-6 related to Alcohols, C12-14-secondary, ethoxylated
Chemical representation	 $n+m = 8 \text{ to } 12$ $x=9$
Molecular formula	CH ₃ (CH ₂) _n CH(CH ₂) _m CH ₂ O(CH ₂ CH ₂ O) _x H, where $n+m=8$ to 12 , and $x=9$
Type of Substance	Fatty alcohol ethoxylate; alkyl PEG ether; oligomer
Appearance	Clear to yellow
Degree of purity	>98%
List of potential impurities with the related concentration (% w/w)	Acetaldehyde ≤ 0.0005% Dioxane ≤ 0.001% Ethyleneglycol ≤ 0.05% Ethyleneoxide ≤ 0.0001%

*Alcohols, C11-15-secondary, ethoxylates is a broad group of substances. Therefore, it should be noted that several trade names can be found under the given CAS No. due to structural differences that can be observed in the final product, e.g., number of ethylene oxide, percentage of homologues with different ranging of n and m .

3. Hazard Identification

3.1. Classification of the substance according to Regulation (EC) No 1272/2008

Acute Toxicity – Category 4 – Oral – H302

Skin Irritation – Category 2 – H315

Serious Eye Damage – Category 1 – H318

Aquatic Chronic – Category 3 – H412

3.2. Classification according to UN GHS

Acute Toxicity – Category 4 – Oral – H302

Acute Toxicity – Category 5 – Dermal – H313

Skin Irritation – Category 2 – H315
Serious Eye Damage – Category 1 – H318
Aquatic Acute – Category 2 – H401
Aquatic Chronic – Category 3 – H412

3.3. Label elements

Pictogram(s) according to UN GHS and Regulation (EC) No 1272/2008



Signal word according to UN GHS and Regulation (EC) No 1272/2008

Danger

Hazard statement(s) according to Regulation (EC) No 1272/2008

H302	Harmful if swallowed
H315	Causes skin irritation
H318	Causes serious eye damage
H412	Harmful to aquatic life with long lasting effects

Hazard statement(s) according to UN GHS

H302	Harmful if swallowed
H313	May be harmful in contact with skin
H315	Causes skin irritation
H318	Causes serious eye damage
H401	Toxic to aquatic life
H412	Harmful to aquatic life with long lasting effects

4. Toxicological information

As mentioned in the introduction, the toxicological evaluation of Deviron® 13-S9 detergent was based on toxicity data collected on members of the alkyl PEG ether group (also called, alcohol ethoxylates) based on a grouping approach. Such clustering was also adopted in formal evaluations conducted by National Authorities, e.g., Australia (IMAP, 2020), Environment Canada (Environment Canada, 2013), and relevant Scientific Committees, e.g., the Cosmetic Ingredient Review (CIR) (Fiume et al., 2012), the EU SCCP (SCCP, 2007).

Acute oral toxicity

Rat, LD₅₀ of 1000 mg/kg based on C₁₄₋₁₅ Pareth 11 (Fiume et al., 2012)

Acute dermal toxicity

Rabbit, LD₅₀ > 2000 mg/kg based on C₁₂₋₁₄ EO₉ (Fiume et al., 2012)

Acute inhalation toxicity

Rat, 8-h exposure to concentrated vapor of PEG-3 methyl ether, no mortality observed; no LC₅₀ established (Fiume et al., 2012)

Skin corrosion/irritation

Based on the collected experimental evidence, it was concluded that C₁₁₋₁₅EO₉ can cause skin irritation.

Serious eye damage/eye irritation

Alcohol ethoxylates were found to be mildly to severely irritating to rabbits' eyes (Fiume et al., 2012). Irreversible effects on the eye, such as corneal opacity, were also reported (Maurer and Kung, 2020).

Respiratory or skin sensitization

Based on the collected evidence, alcohol ethoxylates are not skin sensitizers.

MilliporeSigma, SAFC, Deviron, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.
© 2023 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the US and Canada.

**Millipore
Sigma**

No data available to assess the respiratory sensitization.

Germ cell mutagenicity

Not mutagenic.

Carcinogenicity

Not carcinogenic.

Reproductive and developmental toxicity

Not causing reproduction and developmental toxicity.

NOAEL_{maternal and developmental toxicity}: 50 mg/kg bw/day based on a two-generation rat oral study with C₁₂EO₆. No treatment-related effects on behavior, appearance, survival, or fertility were observed in any of the test groups. No test article-related developmental toxicity effects were observed (Fiume et al., 2012; SCCP, 2007).

NOAEL_{maternal and developmental toxicity}: 50 mg/kg bw/day based on a two-generation rat oral study with C₁₄₋₁₅EO₇. No test-compound related effects on maternal and fetal indices were found. Compound-related effects were limited to increased liver weights (Fiume et al., 2012; SCCP, 2007).

Specific target organ toxicity

Based on available data, repeated exposure is not anticipated to cause significant adverse effects to organs.

NOAEL: 50 mg/kg bw/day based on a 90-day subchronic oral feeding toxicity study in rats with C₁₄₋₁₅EO₇, due to reduced body weight, reduced food intake, increased organ weight, and increased total leukocytes and lymphocytes. However, histopathology revealed no compound-related effects at any dose level. No effects were observed in the organs of the reproductive system. Changes in liver weight, kidney weights and plasma urea concentration were not of toxicological significance (SCCP, 2007).

NOAEL: 50 mg/kg bw/day based on a 2-year chronic dietary feeding study in rats with C₁₄₋₁₅EO₇ or C₁₂₋₁₃EO_{6.5}, due to the increased relative organ weight (Fiume et al., 2012; SCCP, 2007).

Endocrine disrupting properties

Based on an *in silico* (Q)SAR evaluation and on currently available data, no potential for endocrine bioactivity was found in relation to the endpoints evaluated, i.e., estrogen, androgen, and thyroid activity. No data on steroidogenesis were found.

5. Ecological information

The aquatic toxicity of alcohol ethoxylates increases with increasing alkyl chain length and with decreasing EO length (U.S. EPA, 2010), e.g., a C₁₆-alcohol ethoxylate with 3 EO would be more toxic than a C₁₄-alcohol ethoxylate with 9 EO. Furthermore, linear alcohol ethoxylates are more toxic than the branched ones (Environment Canada, 2013). The lowest ecotoxicological values of Deviron® 13-S9 detergent (C₁₁₋₁₅EO₉ branched) were therefore identified among a set of laboratory-based ecotoxicity data collected and assessed on linear and branched alcohol ethoxylates having the same alkyl and ethoxylate chain length of Deviron® 13-S9 detergent.

A QSAR evaluation was also undertaken, where additional ecotoxicity data related to alcohol ethoxylates having a different alkyl chain length and degree of ethoxylation were identified and normalized to C₁₁₋₁₅EO₉, representing Deviron® 13-S9 detergent. Such additional evaluation, which also confirms the proposed GHS classification for aquatic toxicity, is available in the detailed report upon request.

Acute Aquatic Toxicity

Toxicity to fish

96-h mortality, *Pimephales promelas*, LC₅₀ 1.6 mg/L (based C₁₂₋₁₅EO₉ linear) (Dorn et al., 1993)

MilliporeSigma, SAFC, Deviron, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.
© 2023 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the US and Canada.

**Millipore
Sigma**

Toxicity to *Daphnia* and other aquatic invertebrates

48-h immobility, *Daphnia magna*, EC₅₀ 1.3 mg/L (based C₁₂₋₁₅EO₉ linear) (Dorn et al., 1993)

Toxicity to algae

48-h growth, *Raphidocelis subcapitata*, EC₅₀ 4-8 mg/L (based on C₁₂₋₁₄EO₉ linear) (Yamane et al., 1984)

Chronic Aquatic Toxicity

Toxicity to fish

7-d growth, *Pimephales promelas*, NOEC 0.4 mg/L (based C₁₂₋₁₅EO₉ linear) (Dorn et al., 1993)

Toxicity to *Daphnia* and other aquatic invertebrates

7-d growth, *Daphnia magna*, NOEC 1 mg/L (based C₁₂₋₁₅EO₉ linear) (Dorn et al., 1993)

Toxicity to algae

48-h growth (cell density), *Raphidocelis subcapitata*, EC₁₀ 0.151 mg/L (based on C₁₂₋₁₄EO₉ linear) (Yamane et al., 1984 in Belanger et al., 2006)

6. Persistence and degradability

Biodegradability

OECD TG 301B: ≥ 60 % - Readily biodegradable

7. Bioaccumulative potential

- Alcohol ethoxylates are highly metabolizable in fish (Bragin et al., 2020)
- Estimated LogK_{ow} of 4.97 for C₁₁₋₁₅EO₉ based on the LogK_{ow} equation found in Boeije et al., 2006 for alcohol ethoxylates with EO >0.

8. Mobility in soil

- Estimated logK_d of 3.09 (high solid absorption) for C₁₁₋₁₅EO₉, based on the equation developed by van Compernelle et al., 2006 for predicting logK_d of alcohol ethoxylates.

Abbreviations

CIR – Cosmetic Ingredient Review

EC – European Commission

ECHA – European CHemicals Agency

EC₁₀ – concentration causing effects in 10% of a population

EC₅₀ – concentration causing effects in 50% of a population

EO - ethylene oxide

EU – European Union

GHS – Globally Harmonized System of Classification and Labelling of Chemicals

IMAP - Inventory Multi-tiered Assessment and Prioritisation

LC₅₀ – Lethal Concentration causing 50% of deaths

K_d – soil adsorption coefficient

K_{ow} - octanol-water partition coefficient

LD₅₀ – Lethal Dose for 50% of organisms

NOAEL – No Observed Adverse Effect Level

NOEC – No Observed Effect Concentration

OECD – Organisation for Economic Co-operation and Development

PEG - PolyEthylene Glycol

(Q)SAR – (Quantitative) Structure Activity Relationship

REACH - Registration, Evaluation, Authorisation and Restriction of Chemicals

SCCP - Scientific Committee on Consumer Products

TG – Test Guideline

UN – United Nations

U.S. EPA - United States Environmental Protection Agency

References

- Belanger SE, Dorn PB, Toy R, Boeije G, Marshall SJ, Wind T, van Compernelle R, Zeller D. 2006. Aquatic risk assessment of alcohol ethoxylates in North America and Europe. Ecotoxicology and Environmental Safety 64 (2006)

MilliporeSigma, SAFC, Deviron, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2023 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the US and Canada.

**Millipore
Sigma**

85–99.

- Boeije GM, Cano ML, Marshall SJ, Belanger SE, Van Compernelle R, Dorn PB, Gümbel H, Toy R, Wind T. 2006. Ecotoxicity quantitative structure–activity relationships for alcohol ethoxylate mixtures based on substance-specific toxicity predictions. *Ecotoxicology and Environmental Safety* 64 (2006) 75–84.
- Bragin GE, Davis GW, Kung MH, Kelley BA, Sutherland CA, Lampi MA. 2020. Biodegradation and Ecotoxicity of Branched Alcohol Ethoxylates: Application of the Target Lipid Model and Implications for Environmental Classification. *J Surfact Deterg* (2020) 23: 383–403.
- Dorn PB, Salanitro JP, Evans SH, Kravetz L. 1993. Assessing the aquatic hazard of some branched and linear nonionic surfactants by biodegradation and toxicity. *Environmental Toxicology and Chemistry*, 12:1751–1762.
- Environment Canada, 2013. Canadian Environmental Protection Act, 1999. Federal Environmental Quality Guidelines. Alcohol Ethoxylates. Environment Canada. February 2013.
- ECHA, 2023. Guidance for monomers and polymers. February 2023. Version 3.0. Available online at https://echa.europa.eu/documents/10162/23036412/polymers_en.pdf/9a74545f-05be-4e10-8555-4d7cf051bbcd
- Fiume MM, Heldreth B, Bergfeld WF, Belsito DV, Hill RA, Klaassen CD, Liebler D, Marks JG Jr, Shank RC, Slaga TJ, Snyder PW, Andersen AF. 2012. *International Journal of Toxicology* 31(Supplement 2) 169S–244S.
- IMAP (Accelerated Assessment of Industrial Chemicals in Australia). 2020. Ethoxylates of aliphatic alcohols (>C6): Human health tier II assessment. IMAP Group Assessment Report 28/06/2020.
- Maurer LL, Kung MH. 2020. Mammalian Toxicity Testing of Semilinear and Branched Alcohol Ethoxylates. *J Surfact Deterg* (2020) 23: 921–935.
- SCCP, 2007. European Commission, Health & Consumer Protection, Directorate-General. Scientific Committee on Consumer Products (SCCP). Opinion on Polidocanol (Laureth-9). Adopted on 2 October 2007.
- U.S. EPA. 2010. Office of Pollution Prevention and Toxics, 2010. TSCA New Chemical Program (NCP) Chemical Categories.
- van Compernelle R, McAvoy DC, Sherren A, Wind T, Cano ML, Belanger SE, Dorn PB, Kerr KM. 2006. Predicting the sorption of fatty alcohols and alcohols ethoxylates to effluent and receiving water solids. *Ecotoxicology and Environmental Safety* 64 (2006) 61–74.
- Yamane AN, Okada M, Sudo R. 1984. The growth inhibition of planktonic algae due to surfactants used in washing agents. *Water Res.*, 18(9): 1101–1105.

FOR INFORMATIONAL PURPOSES ONLY. We have compiled the content of this document in accordance with our current state of knowledge and legal permissions. The information presented in this document is made available solely for general information purposes. We give no warranty, either expressly or implicitly, for the completeness or correctness of the details provided in this document. Please be aware that the details we provide may no longer be up to date. Access to, use of, and reliance upon this document, any sites via links, and/or any such data, information, or other materials shall be at your own risk. We assume no liability for any inaccuracy of this document or its contents. We reserve the right to withdraw or amend this document, and any products or service we provide, at any time and in our sole discretion, without notice. This document and its contents are provided on an “as is” and “as available” basis, without any warranties of any kind, either express or implied. Neither we nor any person associated with us make any express or implied representations or warranties regarding the usability of this document (including, without limitation, any data, information, or other materials), or represent or endorse the accuracy, suitability, reliability, availability, and/or completeness of any data, information, or other materials in any manner in this document.

PROPRIETARY. Unless expressly stated otherwise, this document and the information contained within, including the document structure, contents, and its attachments, are the property of Merck KGaA, Darmstadt, Germany and/or its affiliates and are protected by copyright laws.

MilliporeSigma, SAFC, Deviron, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.
© 2023 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the US and Canada.

**Millipore
Sigma**