

Technical Bulletin

Trypsin inhibitor from Glycine max (soybean)

Solution, sterile-filtered, suitable for, suitable for cell culture

T6414

Storage Temperature -20 °C

Product Description

CAS Number

9035-81-8

Extinction Coefficient

$E^{1\%} = 9.94$ (280 nm, pH 7.6 buffer)

pI

4.5¹

Synonyms

Kunitz Trypsin Inhibitor, Ti^{a1}, STI, and SBT1

This product is a 1% solution of trypsin inhibitor in Dulbecco's PBS without calcium and magnesium. There is also added a small amount (2% v/v) of charcoal stripped FBS to stabilize the cell membrane. This product does contain some α_1 antitrypsin activity.

This product is cell culture tested and is appropriate for use in cell culture applications. It has been optimized for passage of endothelial cell cultures.

Soybean trypsin inhibitor was first isolated by Kunitz.² Several other related inhibitors are also found in soybeans.³ Trypsin inhibitor from soybeans is a monomeric protein containing 181 amino acid residues in a single polypeptide chain crosslinked by two disulfide bridges.^{4,5,6} The molecular weight determined from the amino acid sequence is 20.1 kDa.

Soybean trypsin inhibitor inhibits trypsin, and to a lesser extent chymotrypsin⁸ and plasmin.⁹ Soybean trypsin inhibitor will also inhibit proteases with mechanisms similar to trypsin. This product will also inhibit plasma kallikrein and coagulation Factor X_a. However, this product will not inhibit metalloproteases, tissue-based kallikrein, acid proteases, or thio proteases.

Soybean trypsin inhibitor forms a 1:1 stoichiometric complex with the protease active site. Upon formation of this complex, trypsin may cleave a single arginine- isoleucine bond on the inhibitor.^{10,11} Inhibition is both reversible and pH dependent. Dissociation of this complex may yield a modified or native form of the inhibitor.¹² The optimal pH for trypsin binding is 8.0 with an association constant of greater than 109 at pH 8.0 and $0.15\text{--}2.6 \times 10^4$ at pH 3.6 to 4.4.¹³

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses.

Storage/Stability

This product is stable in frozen aliquots at -20 °C, but freeze-thaw should be avoided. This protein is reversibly denatured by short heating to 80 °C and irreversibly inhibited by heating to 90 °C.³

Procedure

After trypsinizing cells, resuspend cells in 1 mL trypsin inhibitor solution (1 mg/mL) for every mL of trypsin solution used for dissociation. Centrifuge the cell suspension at 1000 rpm (200 x g) for 5 minutes. A cell pellet should form. Remove as much of the trypsin inhibitor as possible and resuspend the pellet in serum-free medium. Culture cells as desired.

References

1. Kunitz, M., J. Gen. Physiol., 30, 311 (1947).
2. Kunitz, M., Science, 101, 668 (1945).
3. Steiner, R. F., and Frattali, V. J., Agr. Food Chem., 17, 513 (1969).
4. Kim, S. H., et al., Comparative study on amino acid sequences of Kunitz-type soybean trypsin inhibitors, Tia, Tib, and Tic. J. Biochem. (Tokyo), 98(2), 435- 448 (1985).
5. Steiner, R. F., Biochim. Biophys. Acta, 100, 111 (1965).
6. Koide, T., and Ikenaka, T., Studies on soybean trypsin inhibitors. 1. Fragmentation of soybean trypsin inhibitor (Kunitz) by limited proteolysis and by chemical cleavage. Eur. J. Biochem., 32(3), 401-407 (1973).
7. Wu, Y. V., and Scheraga, H. A., Biochemistry, 1, 698 (1962).
8. De Vonis Bidlingmeyer, U., et al., Identity of the tryptic and alpha-chymotryptic reactive sites on soybean trypsin inhibitor (Kunitz). Biochemistry, 11(17), 3303-3310 (1972).
9. Naninga, L. B., and Guest, M. M., Arch, Biochem. Biophys., 108, 542 (1964).
10. Ozawa, K., and Laskowski, M. Jr., The reactive site of trypsin inhibitors. J. Biol. Chem., 241(17), 3955-3961 (1966).
11. Finkenshtadt, W. R., and Laskowski, M. Jr., Resynthesis by trypsin of the cleaved peptide bond in modified soybean trypsin inhibitor. J. Biol. Chem., 242(4), 771-773 (1967).
12. Finkenshtadt, W. R., and Laskowski, M. Jr., J. Biol. Chem., 240, 962 (1965).
13. Advances in Protein Chemistry, Vol. 9, Laskowski, M., Academic Press (New York, NY: 1954), p. 226.

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at [SigmaAldrich.com/techservice](https://www.sigmaaldrich.com/techservice).

Standard Warranty

The applicable warranty for the products listed in this publication may be found at [SigmaAldrich.com/terms](https://www.sigmaaldrich.com/terms).

Contact Information

For the location of the office nearest you, go to [SigmaAldrich.com/offices](https://www.sigmaaldrich.com/offices).

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.

Merck and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2021 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

T6414 Rev 10/21

The Merck logo, consisting of the word "MERCK" in a bold, red, sans-serif font.