

Product Information

Deoxyribonuclease I from bovine pancreas

Sigma Type IV, lyophilized powder, ≥ 2000 Kunitz units/mg protein**D5025**

Product Description

CAS Registry Number: 9003-98-9

Enzyme Commission (EC) Number: 3.1.21.1

Synonyms: DNase I, Deoxyribonuclease A, Deoxyribonuclease 5'-oligonucleotidohydrolase

Deoxyribonuclease I (DNase I) is an endonuclease that cleaves DNA by preferentially acting on phosphodiester bonds adjacent to pyrimidines, to produce polynucleotides with terminal 5'-phosphates. A tetranucleotide is the smallest average digestion product. In the presence of Mg^{2+} ions, DNase I attacks each strand of DNA independently and the cleavage sites are random. If Mn^{2+} ions are present, both DNA strands are cleaved at approximately the same site.¹ DNase I hydrolyzes single-stranded DNA, double-stranded DNA, and chromatin (the reaction rate is restricted by DNA association with histones).

DNase I is found in most cells and tissues. In mammals, the pancreas is one of the best sources for the enzyme. Pancreatic DNase I was the first DNase to be isolated. The calculated molecular mass is 30,072 Da. DNase I exists as a mixture of glycoproteins with two disulfide bridges.²

Bovine pancreatic DNase I contains four chromatographically distinguishable components, labeled A, B, C, and D.³ The molar ratios of A:B:C in a pancreatic extract are 4:1:1. Only minor amounts of D are found. Forms A and B differ in carbohydrate content (see Table below).⁴

Carbohydrate Content⁴

Carbohydrate / Form	A	B	C
N-Acetylglucosamine	2	3	2
Mannose	6	5	5
Sialic Acid	-	1	-
Galactose	-	1	-

Form C differs from Forms A and B by having one less His and one more Pro, and in the carbohydrate chain.⁴

DNase I is used to remove DNA from protein and nucleic acid samples, and to nick DNA as a first step to incorporate labeled bases into DNA. Several theses⁵⁻⁷ and dissertations⁸⁻²⁶ have cited use of product D5025 in their protocols.

Isoelectric points:²

- A: 5.22
- B: 4.96
- C: 5.06
- D: 4.78

Optimal pH: 7-8

Extinction Coefficient: $E_{280}^{1\%} = 11.1$

Activators

- DNase I has an absolute requirement for divalent metal cations.
- The most commonly used divalent metal cation is Mg^{2+} .^{27,28}
- However, Mn^{2+} , Ca^{2+} , Co^{2+} , and Zn^{2+} will activate DNase I.²⁷⁻²⁹
- 5 mM Ca^{+2} will stabilize DNase I against proteolytic digestion.³⁰
- 0.1 mM Ca^{+2} is needed to reduce the rate of inactivation by one-half.³⁰

Inhibitors

There is no general inhibitor specific for DNase I.^{27,28} Citrate inhibits Mg^{2+} -activated DNase I, but not Mn^{2+} -activated DNase I.

- 2-Mercaptoethanol (the reduced enzyme is inactive, but can be reactivated in the presence of Ca^{2+} or Mg^{2+} ions)²⁸
- Chelators (such as EDTA, EGTA)
- Sodium dodecyl sulfate (SDS)³⁰
- Actin³²

MERCK

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Product

This product is purified from bovine pancreas. The purification procedure is not selective for any form (A, B, C, or D) of DNase I. D5025 is supplied as a lyophilized powder, containing CaCl_2 .

Protein content: $\geq 80\%$

Specific activity: ≥ 2000 Kunitz units/mg protein

Unit Definition:³³

- One Kunitz unit will produce a ΔA_{260} of 0.001 per minute per mL at pH 5.0 at 25 °C, using DNA, Type I or III, as substrate, with $[\text{Mg}^{2+}] = 4.2 \text{ mM}$.
- This enzyme assay reaction is performed in 95 mM acetate buffer, pH 5.0, at 25 °C, containing 4.75 mM Mg^{2+} and 1.9 mM Ca^{2+} , in a 3 mL reaction.

Contaminants:

- Protease: $\leq 0.05\%$
- RNase: $\leq 0.02\%$
- Chymotrypsin: $\leq 0.5\%$

Preparation Instructions

This product is tested for solubility in 0.15 M NaCl at 5 mg/mL. One publication has reported preparation of stock solutions of this DNase I product at 20 mg/mL in Ringer's solution,³⁴ although we have not tested this condition ourselves.

Storage/Stability

D5025 has a recommended retest date of three years, when unopened and stored long-term at the recommended temperature, -20°C .

Solutions of DNase I at 10 mg/mL in 0.15 M NaCl may lose $<10\%$ of its activity when stored for a week in aliquots at -20°C . The same solutions stored in aliquots at $2-8^\circ\text{C}$ can lose $\sim 20\%$ activity.

DNase I remains active in solution between pH 5-7 up to 60°C for at least five hours. A 1 mg/mL solution in acetate buffer (pH 5.0) or Tris buffer (pH 7.2) loses activity at the rate of 6% per hour. At 68°C , DNase I loses activity in <10 minutes.

References

1. Sambrook, J. *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd edition. Cold Spring Harbor Laboratory Press (Cold Spring Harbor, NY), Vol. 2, p. 5.83 (1989).
2. Liao, T.-H. *et al.*, *J. Biol. Chem.*, **248(4)**, 1489-1495 (1973).
3. Kim, H.S., and Liao, T.-H., *Anal. Biochem.*, **119(1)**, 96-101 (1982).
4. Salnikow, J. *et al.*, *J. Biol. Chem.*, **245(21)**, 5685-5690 (1970).
5. Das, Madhurima, "Studies of Regulated Exocytosis from Neuroendocrine Cells". Wayne State University, M.S. thesis, p. 33 (2013).
6. Bacon, Margaret Leese, "Identification of Clathrin and Dynamin II in the Porcine Ovary Supports the Presence of Clathrin-Mediated Endocytosis". Colorado State University, M.S. thesis, p. 34 (2016).
7. West, Michael, "Enhanced Vascularization of Modular Engineered Tissues Using Activated Macrophages". University of Toronto, M.Sc. thesis, p. 16 (2016).
8. Davis, Joe William, "Intracellular Degradation of Ribosomes in a Mesophilic and a Thermophilic Species of *Bacillus*". Western Michigan University, Ph.D. dissertation, p. 14 (1990).
9. Jacobson, Derek William Niall, "Imaging the mitochondrial permeability transition in single cells: free radicals and calcium as determinants of cell fate". University College London, Ph.D. dissertation, p. 24 (2000).
10. Mouzannar, Raymond, "Higher order chromatin degradation induced by hydrogen peroxide in glial cells". West Virginia University, Ph.D. dissertation, p. 56 (2001).
11. Kaesberg, Jeffrey, "Bacteriophages of *Xanthomonas Campestris* P.v. *Begoniae*; Their Occurrence, Survival and Potential Use as a Biological Control Agent". University of Florida, Ph.D. dissertation, p. 33 (2009).
12. Nguyen, Tina T., "Structural and Functional Studies of Proteins Involved in Antigen Processing". University of Massachusetts Worcester, Ph.D. dissertation, p. 183 (2010).
13. Fedele, Vita, "Polycomb proteins and breast cancer". University of St. Andrews, Ph.D. dissertation, p. 69 (2012).
14. Hernández, Natalia M. Ramos, "The Role of NDFIP1 in T Cell Tolerance". University of Pennsylvania, Ph.D. dissertation, pp. 37, 79 (2013).

15. Eucker, Tyson Paul, "Campylobacter jejuni Manipulates the Focal Complex to Invade Epithelial Cells and Induce Secretion of Interleukin-8". Washington State University, Ph.D. dissertation, p. 37 (2014).
16. Green, Ryan Liam, "Investigating the role of FUS, TDP-43 and DYNC1H1 mutations in the etiology of adult and childhood-onset motor neuron disease". University of Sussex, Ph.D. dissertation, pp. 93, 96 (2016).
17. Katic, Jelena, "The role of the cell adhesion molecule CHL1 and its interaction partners in cerebellar development". Universität Hamburg, Dr. rer. nat. dissertation, p. 42 (2016).
18. Capitanio, Juliana Silva, "Nup98-mediated Regulation of Gene Expression and RNA Metabolism". University of Alberta, Ph.D. dissertation, p. 76 (2017).
19. D'Innocenzi, Jaclyn Marie, "Non-Small Cell Lung Cancer utilizes SIRT6 to initiate tumorigenesis and overcome cellular senescence". University of Virginia, Ph.D. dissertation, p. 53 (2017).
20. Kim, Eun Joo, "Spatiotemporal Effects of Dietary Bioactives on LGR5⁺ Stem Cells During Colon Tumorigenesis". Texas A&M University, Ph.D. dissertation, p. 269 (2017).
21. Eisel, David, "Reprogramming of M2-like macrophages by CD4⁺ T cells, transcription factor knockdown and miRNA transfection". Ruperto-Carola University Heidelberg, Dr. rer. nat. dissertation, p. 34 (2018).
22. Kim, Jee Soo (Monica), "Development of a high-throughput drug screening platform for oligodendrocyte myelination (for progressive multiple sclerosis)". University of Edinburgh and University of Strathclyde, Ph.D. dissertation, p. 99 (2019).
23. Nabhan, Ahmad N., "Alveolar stem cells and niches in aging, injury and evolution". Stanford University, Ph.D. dissertation, p. 91 (2019).
24. Geller, Anne Elena, "Harnessing the power of trained immunity in the setting of pancreatic cancer: a novel mechanism of immune trafficking and tumor control". University of Louisville School of Medicine, Ph.D. dissertation, p. 167 (2021).
25. Follis, Rose M., "Metabolic control of sensory neuron survival by the p75 neurotrophin receptor in Schwann cells". Vanderbilt University, Ph.D. dissertation, pp. 57, 118 (2021).
26. van der Gucht, Marie, "Understanding and Exploiting the Phage Toolbox for Biomedical and Biotechnological Applications". KU Leuven, Ph.D. dissertation, p. 35 (2021).
27. Weir, A.F., *Methods Mol. Biol.*, **16**, 1-6 (1993).
28. Weir, A.F., *Methods Mol. Biol.*, **16**, 7-16 (1993).
29. Price, P.A. et al., *J. Biol. Chem.*, **244(4)**, 929-932 (1969).
30. Price, P.A. et al., *J. Biol. Chem.*, **244(4)**, 917-923 (1969).
31. Liao, T.-H., *J. Biol. Chem.*, **250(10)**, 3831-3836 (1975).
32. Lazarides, E., and Lindberg, U., *Proc. Natl. Acad. Sci. USA*, **71(12)**, 4742-4746 (1974).
33. Kunitz, M., *J. Gen. Physiol.*, **33(4)**, 349-362 (1950).
34. Baumann, T.K., "Cultures of Adult Trigeminal Ganglion Neurons", in *Methods in Toxicology, Volume 1: Part A – In Vitro Biological Systems* (C.A. Tyson and J.M. Frazier, eds.). Academic Press (San Diego, CA), pp. 61-69 (1993).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Standard Warranty

The applicable warranty for the products listed in this publication may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.

Merck and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2022 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.
D5025pis Rev 06/22 VLR,JWM,RGB,GCY,MAM

MERCK