3050 Spruce Street, St. Louis, MO 63103 USA
Tel: (800) 521-8956 (314) 771-5765 Fax: (800) 325-5052 (314) 771-5757
email: techservice@sial.com sigma-aldrich.com

Product Information

Anti-Mouse IgG (Fab specific)

produced in goat, affinity isolated antibody adsorbed with human IgG

Catalog Number M6898

Product Description

Anti-mouse IgG (Fab specific) is produced in goat using purified mouse IgG, Fab fragment, as the immunogen. Antibody is isolated from goat anti-mouse IgG antiserum by immunospecific purification which removes essentially all goat serum proteins, including immunoglobulins, which do not specifically bind to the Fab fragment of mouse IgG. The antibody preparation is solid phase adsorbed with human IgG to ensure minimal cross reactivity in tissue or cell preparations.

Specificity is determined by immunoelectrophoresis (IEP). By IEP single precipitation arcs are observed against normal mouse serum, mouse IgG and the Fab fragment of mouse IgG, while no reaction is observed against the Fc fragment of mouse IgG. No reaction is observed with human IgG.

Identity and purity of the antibody is established by immunoelectrophoresis. Electrophoresis of the antibody preparation, followed by diffusion versus anti-goat IgG and anti-goat whole serum, results in single arcs of precipitation.

Reagent

Supplied as a solution in 0.01 M phosphate buffered saline, pH 7.4, containing 15 mM sodium azide as a preservative.

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

Store at 2-8 °C for up to one month. For extended storage, the solution may be frozen in working aliquots. Repeated freezing and thawing, or storage in "frost-free" freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use.

Product Profile

Indirect ELISA: a working dilution of 1:20,000 was determined using 5 µg/ml mouse IgG for coating

Note: In order to obtain the best results in various techniques and preparations we recommend determining the optimal working dilutions by titration.

DS,KAA,PHC 05/13-1