

## Product Information

# Nystatin

Powder, suitable for cell culture, BioReagent

**N6261**

Store at 2-8 °C.

## Product Description

**Molecular Formula:** C<sub>47</sub>H<sub>75</sub>NO<sub>17</sub>

**Molecular Weight:** 926.1

**CAS Number:** 1400-61-9

**Melting Point:** Gradually decomposes above 160 °C without melting by 250 °C

**λ<sub>max</sub>:** 290, 307, and 322 nm (ethanol)

**Specific Rotation (at 25 °C):**

-10° (glacial acetic acid)

+25° (pyridine)

+12° (DMF)

-7° (0.1 N HCl in methanol)

This product is cell culture tested and recommended for use in cell culture applications (50 mg/L). This is a non-sterile powder and is not recommended for aseptic work.

Nystatin is a fungistatic and fungicidal polyene antibiotic, which increases the permeability of the cell membrane of sensitive fungi by binding to sterols, chiefly ergosterol.<sup>1</sup> Its main action is against *Candida* species. It is also effective against *Aspergillus*, *Coccidioides immitis*, *Cryptococcus neoformans*, *Histoplasma capsulatum*, *Blastomyces dermatidis*, and other yeasts and fungi.<sup>2</sup> Nystatin has been used to enrich mutants by killing yeast cells.<sup>5</sup> Nystatin has no antibacterial activity.<sup>4</sup> The minimum inhibitory concentration for most sensitive fungi has been reported to range from 1.56 to 6.25 µg/mL.

Nystatin is poorly absorbed from the gastrointestinal tract. It is not absorbed through the skin or mucous membranes when applied topically.

## Precautions and Disclaimer

For Laboratory Use Only. Not for drug, household or other uses.

## Preparation Instructions

Nystatin is soluble at 28 °C in the following: methanol (11.2 mg/mL), ethanol (1.2 mg/mL), carbon tetrachloride (1.23 mg/mL), chloroform (0.48 mg/mL), benzene (0.28 mg/mL), and ethylene glycol (8.75 mg/mL).

Nystatin is soluble in DMSO (5 mg/mL), yielding a clear, bright yellow solution. This product is freely soluble in DMF and formamide.<sup>3</sup> A 3% suspension in water has pH 6.5-8.0.<sup>2</sup>

It is not recommended to autoclave or sterile filter solutions of Nystatin. If sterility is required, use γ-irradiated product N4014.

## Storage/Stability

Solutions and aqueous suspensions begin to lose activity soon after preparation. Heat, light, and oxygen accelerate decomposition. Aqueous suspensions are stable for 10 minutes when heating to 100 °C at pH 7. Nystatin (in tissue culture media) is stable at 37 °C for three days. It is also stable in moderately alkaline media, but labile at pH 9 and 2. Activity is not diminished by blood or serum.<sup>1</sup>

Nystatin, cell culture tested (N1638) is a sterile nystatin suspension in Dulbecco's Phosphate Buffered Saline. This product is stored frozen and shipped on dry ice. When stored frozen it has a shelf life of 24 months.

## References

1. Brezis, M., Rosen, S., Silva, P., Spokes, K., & Epstein, F. H. (1984). Polyene toxicity in renal medulla: injury mediated by transport activity. *Science*, 224(4644), 66-68.
2. Reynolds, J. E. (Ed.). (1989). *Martindale: the extra pharmacopoeia*, pp. 432-433.
3. Moffat, A. C. (1986). Clarke's Isolation and Identification of Drugs in Pharmaceuticals. *Body Fluids and Post-Mortem Material.*, p. 829.
4. Upson, D. W. (1985). *Upson's handbook of clinical veterinary pharmacology* (No. Edition 2, p. 660pp)., p. 640.
5. Hartwell, L. H. (1970). Biochemical genetics of yeast. *Annual review of genetics*, 4(1), 373-396.

## Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

### Technical Assistance

Visit the tech service page at [SigmaAldrich.com/techservice](https://SigmaAldrich.com/techservice).

### Terms and Conditions of Sale

Warranty, use restrictions, and other conditions of sale may be found at [SigmaAldrich.com/terms](https://SigmaAldrich.com/terms).

### Contact Information

For the location of the office nearest you, go to [SigmaAldrich.com/offices](https://SigmaAldrich.com/offices).

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the U.S. and Canada.

MilliporeSigma and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2008-2025 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

23240377 Rev 07/25

**MILLIPORE  
SIGMA**