3050 Spruce Street, St. Louis, MO 63103 USA Tel: (800) 521-8956 (314) 771-5765 Fax: (800) 325-5052 (314) 771-5757 email: techservice@sial.com sigma-aldrich.com

Product Information

G 418 disulfate salt

Product Codes **G 5013**, **G 8168**, **A 1720**, **G 1279**, **and A 8601**Storage Temperature 2-8 °C

CAS# 108321-42-2

Product DescriptionMolecular weight: 692.7

Molecular formula: $C_{20}H_{40}N_4O_{10} \bullet 2H_2SO_4$ Specific Rotation: +104.4° (c = 0.3% in H_2O at 26 °C)

G 418 is an aminoglycoside antibiotic similar in structure to gentamycin. It exhibits toxicity towards both eukaryotic and prokaryotic cells. The optimal concentration for selection and maintenance must be determined for each cell line. For bacteria and algae, concentrations of 5 $\mu g/ml$ or less are recommended. Animal cells may require up to 300-500 $\mu g/ml$. Typically, resistance is conferred by one of two dominant genes of bacterial origin, which can be expressed in eukaryotic cells. Cells that are multiplying will be affected sooner than those that are not. Cells in log phase may require three to seven days for selection. In general, concentrations of approximately 400 $\mu g/ml$ for selection and 200 $\mu g/ml$ for maintenance are required for mammalian cells.

Reagents

Products G 5013, A 1720, G 1279, and A 8601 are sold as powders. G 8168 is a 5% aqueous solution prepared from A 1720.

Products A 1720 and G 8168 are cell culture tested. Product G 1279 is tested for plant cell culture applications. A 8601 is Biotechnology Performance Certified grade.

Preparation Instructions

The G 418 powder is soluble in water at 100 mg/ml.

Storage/Stability

The G 418 powder is stable for three years as supplied when stored at 2-8 °C. The G 418 solution (Product Code G 8168) is stable for two years at 2-8 °C

References

- Loebenberg, D., et al., G 418, a new micrommomospora-produced aminglycoside with activity apainst protozoa and helminths: antiparasitic activity. Antimicrob. Agents Chemother., 7, 811 (1975).
- Ursic, D., et al., A new antibiotic with known resistance factors, G 418, inhibits plant cells. Biochem. Biophys. Res. Commun., 101, 1031 (1981).
- 3. Colbere-Garapin, F., et al., A new dominant hybrid selective marker for higher eukaryotic cells. J. Mol. Biol., **150**, 1 (1981).
- 4. Jimenez, A., and Davies, J., Expression of a transposable antibiotic resistance element in *Saccharomyces*. Nature, **287**, 869 (1980).
- Hirth, K.P., et al., A DNA-mediated transformation system for *Dictyostelium discoideum*. Proc. Natl. Acad. Sci. USA, 79, 7356 (1982).

LMY, MAM, SBC 03/21-1