

## Product Information

# Cytochrome c from bovine heart

**C2037**

Storage Temperature -20 °C.

CAS# 9007-43-6

Synonym: Ferricytochrome c (oxidized state)

## Product Description

Cytochrome c is an electron-carrying mitochondrial protein composed of 104 amino acid residues.<sup>1</sup> It is a heme protein containing a single polypeptide chain and a single heme group, which is covalently attached to the polypeptide. The ready fluctuation of cytochrome c within the cell between the ferrous and ferric states, makes it an efficient biological electron-transporter and it plays a vital role in cellular oxidations in both plants and animals. It is generally regarded as a universal catalyst of respiration, forming an essential electron-bridge between the respirable substrates and oxygen.

This Cytochrome c product is prepared from bovine heart by a modification of a published method.<sup>2</sup> The product is supplied as a lyophilized powder. The final step before lyophilization is extensive dialysis against 6 mM Ammonium hydroxide, which is volatile under lyophilization conditions, so the final product should not contain any buffer salts.

The product is mainly the oxidized form of the protein. The reduced form of cytochrome c can be prepared with either Sodium dithionite or Sodium ascorbate, followed by gel filtration.<sup>3</sup>

Molecular mass:<sup>4</sup> 12,327 g/mol

Isoelectric point (pI):<sup>5</sup> range of 10.0–10.5

Spectral properties:<sup>6</sup>

- $\lambda_{\max}$  = 550 nm (reduced form)
- $E_{\text{mM}}^{\text{M}}$  = 28.0 (reduced form, 0.1 M phosphate buffer, pH 6.8)
- $E_{\text{mM}}^{\text{M}}$  = 8.4 (oxidized form, 0.1 M phosphate buffer, pH 6.8)
- Purity:  $\geq 95\%$  (SDS-PAGE and spectral assay)

## Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

## Preparation Instructions

Cytochrome c is soluble in water or buffered solutions at pH  $7.0 \pm 1.0$  (up to 200 mg/mL), yielding a clear, dark red solution. For general purposes reconstitution is done at 10 mg/mL.

## Storage/Stability

Store cytochrome c at -20 °C. The product, as supplied, is stable for 5 years.

### Recommended storage time for aqueous solutions

- Storage at -20 °C (freezer) – 6 months
- Storage at 2-8 °C (refrigerator) – 2 weeks
- Storage at 20-25 °C (ambient temperature) – 3 days

## References

1. Nakashima, T. et al., J. Biol. Chem., 241(5), 1166- 1177 (1966).
2. Hagihara, I. et al., Biochem. Prep., 6, 1 (1958).
3. Dixon, H.B., and McIntosh, R., Nature, 213(74), 399-400 (1967).
4. Handbook of Biochemistry (CRC Press, 1968).
5. Brain Research, 153, 477-493 (1978).
6. Van Gelder, B.F., and Slater, E.C., Biochem. Biophys. Acta, 58, 593-595 (1962).

## Notice

---

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

### Technical Assistance

Visit the tech service page at [SigmaAldrich.com/techservice](https://SigmaAldrich.com/techservice).

### Terms and Conditions of Sale

Warranty, use restrictions, and other conditions of sale may be found at [SigmaAldrich.com/terms](https://SigmaAldrich.com/terms).

### Contact Information

For the location of the office nearest you, go to [SigmaAldrich.com/offices](https://SigmaAldrich.com/offices).

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the U.S. and Canada.

The vibrant M, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.  
© 2024 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

Document # Rev MM/YY

